



12-1998

Translational Shape Invariance and the Inherent Potential Algebra

Asim Gangopadhyaya

Loyola University Chicago, agangop@luc.edu

Jeffrey Mallow

Loyola University Chicago, jmallow@luc.edu

Uday P. Sukhatme

University of Illinois at Chicago, sukhatme@uic.edu

Follow this and additional works at: https://ecommons.luc.edu/physics_facpubs



Part of the [Physics Commons](#)

Recommended Citation

Gangopadhyaya, Asim, Jeffrey V. Mallow, and Uday P. Sukhatme. "Translational Shape Invariance and the Inherent Potential Algebra." *Physical Review A* 58, no. 6 (December 1, 1998): 4287–92. doi:10.1103/PhysRevA.58.4287.

This Article is brought to you for free and open access by the Faculty Publications and Other Works by Department at Loyola eCommons. It has been accepted for inclusion in Physics: Faculty Publications and Other Works by an authorized administrator of Loyola eCommons. For more information, please contact ecommons@luc.edu.



This work is licensed under a [Creative Commons Attribution-NonCommercial-No Derivative Works 3.0 License](#).
© 1999 The American Physical Society.

Translational shape invariance and the inherent potential algebra

Asim Gangopadhyaya,^{1,*} Jeffrey V. Mallow,^{1,†} and Uday P. Sukhatme^{2,‡}

¹*Department of Physics, Loyola University Chicago, Chicago, Illinois 60626*

²*Department of Physics, University of Illinois at Chicago (m/c 273), 845 West Taylor Street, Chicago, Illinois 60607-7059*

(Received 8 May 1998)

For all quantum-mechanical potentials that are known to be exactly solvable, there are two different, and seemingly independent methods of solution. The first approach is the potential algebra of symmetry groups; the second is supersymmetric quantum mechanics, applied to shape-invariant potentials, which comprise the set of known exactly solvable potentials. Using the underlying algebraic structures of Natanzon potentials, of which the translational shape-invariant potentials are a special subset, we demonstrate the equivalence of the two methods of solution. In addition, we show that, while the algebra for the general Natanzon potential is $so(2,2)$, the subgroup $so(2,1)$ suffices for the shape invariant subset. Finally, we show that the known set of exactly solvable potentials in fact constitutes the full set of such potentials. [S1050-2947(98)09611-5]

PACS number(s): 03.65.Fd, 03.65.Sq, 02.20.-a

I. INTRODUCTION

Natanzon potentials [1] form a complete set of exactly solvable potentials of nonrelativistic quantum mechanics for which the Schrödinger equation reduces to the hypergeometric equation. In Refs. [2,3], Alhassid and co-workers have studied the group structure of these quantum-mechanical systems, related their Hamiltonians to the Casimir operator of an underlying $so(2,2)$ algebra, and determined all their quantum states by group-theoretical methods.

In supersymmetric quantum mechanics (SUSY-QM) [4] one applies a different algebraic method. The exactly solvable problems in SUSY-QM are described by superpotentials $W(x,a)$ that obey a special integrability condition,

$$W^2(x,a_0) + \frac{dW(x,a_0)}{dx} = W^2(x,a_1) - \frac{dW(x,a_1)}{dx} + R(a_0), \quad (1)$$

known as shape invariance [5,6]. $R(a_0)$ is a constant and the parameter a_1 is a function of a_0 ; i.e., $a_1 = f(a_0)$. For a shape-invariant system, the entire spectrum can be determined algebraically by a procedure similar to that of the one-dimensional harmonic oscillator, without ever referring to the underlying differential equations. Although most of the known shape-invariant potentials (SIP) belong to the Natanzon class, there are a few exceptions [7,8].

In a previous work [9], we have shown that problems for which (1) there is a translational change of parameters $a_1 = f(a_0) = a_0 + \text{const}$ and (2) $R(a_0)$ is a linear function of a_0 , the shape-invariance condition of Eq. (1) implies the presence of an $so(2,1)$ dynamical algebra. Hence these problems are solvable by either method. As shown in Ref. [4], potentials of this type include Morse, Scarf I, Scarf II, and

generalized Pöschl-Teller. However, the formalism used in Ref. [9] is not readily extendable to other shape-invariant potentials.

In this paper, we generalize our work of Ref. [9]. However, we have used a different approach here that is closely based on the work of Alhassid, Gürsey, and Iachello [2]. The authors of Ref. [2] have shown that a Hamiltonian with a general Natanzon potential has an $so(2,2)$ symmetry. We study here the algebra of Natanzon potentials that are also shape invariant. We find that general Natanzon potentials when subjected to a further constraint give the entire set of shape-invariant potentials. The shape-invariant potentials that reduce to the confluent hypergeometric equation can be obtained as a limit [10]. We also find that while the algebra for the general Natanzon potential is $so(2,2)$, a subgroup $so(2,1)$ suffices for all the shape-invariant problems of the Natanzon type.

Thus, this paper connects all the shape-invariant potentials of translational type ($a_1 = a_0 + \text{const}$) to an algebraic structure that has many interesting consequences. Some time ago it was discovered that spectra of potentials with translational shape invariance can be exactly determined by the supersymmetric WKB method [11], which usually only gives approximate results. The reason for this exactness was very puzzling. However, in light of this group-theoretical connection, this result may not be that difficult to understand, as various authors have demonstrated the exactness of WKB results on a group manifold [12].

In Sec. II, we will quickly review the formalism of SUSY-QM and shape invariance. In Sec. III, we will briefly describe our previous work [9] where we connected a subset of SIP's to $so(2,1)$ potential algebra. In Sec. IV, we discuss the potential algebra of a general Natanzon potential. This section will closely follow Ref. [2]. In Sec. V, we identify conditions under which the general Natanzon potential reduces to a shape-invariant potential. We then show that this condition has a finite number of solutions for shape-invariant potentials; however, they generate all the known shape-invariant potentials of translational type.

*Electronic address: agangop@luc.edu, asim@uic.edu

†Electronic address: jmallow@luc.edu

‡Electronic address: sukhatme@uic.edu

II. SUSY-QM AND SHAPE INVARIANCE

In this section, we very briefly describe supersymmetric quantum mechanics (SUSY-QM), and also show how SUSY-QM with shape invariance allows one to completely determine the spectrum of a quantum system. For a more detailed description, see Ref. [4].

A quantum-mechanical system given by a potential $V_-(x)$ can alternatively be described by its ground-state wave function $\psi_0^{(-)}$. From the Schrödinger equation for the ground-state wave function $[-\psi_0'' + V(x)\psi_0 = 0]$, it follows that the potential can be written as $V_-(x) = (\psi_0''/\psi_0)$, where the prime denotes differentiation with respect to x . (It is assumed that the potential is properly adjusted to make the ground-state energy $E_0 = 0$.) In SUSY-QM, it is customary to express the system in terms of the superpotential $W(x) = -(\psi_0'/\psi_0)$. The ground-state wave function is then given by $\psi_0 \sim \exp[-\int_{x_0}^x W(x)dx]$; x_0 is an arbitrarily chosen reference point. The Hamiltonian H_- can now be written as

$$\begin{aligned} H_- &= \left(-\frac{d^2}{dx^2} + V_-(x) \right) = \left(-\frac{d^2}{dx^2} + W^2(x) - \frac{dW(x)}{dx} \right) \\ &= \left(-\frac{d}{dx} + W(x) \right) \left(\frac{d}{dx} + W(x) \right). \end{aligned} \quad (2)$$

(We are using units with \hbar and $2m = 1$.) In analogy with the harmonic-oscillator raising and lowering operators, we have introduced two operators: $A = [(d/dx) + W(x)]$, and its Hermitian conjugate $A^\dagger = [-(d/dx) + W(x)]$. Thus $H_- = A^\dagger A$.

However, one can easily construct another Hermitian operator $H_+ = AA^\dagger$ and show that the eigenstates of H_+ are isospectral with excited states of H_- . The Hamiltonian H_+ , with potential $V_+(x) = \{W^2(x) + [dW(x)/dx]\}$, is called the superpartner of the Hamiltonian H_- . To show the isospectrality mentioned above, let us denote the eigenfunctions of H_\pm that correspond to eigenvalues E_n^\pm , by $\psi_n^{(\pm)}$. For $n = 1, 2, \dots$,

$$\begin{aligned} H_+(A\psi_n^{(-)}) &= AA^\dagger(A\psi_n^{(-)}) = A(A^\dagger A\psi_n^{(-)}) \\ &= AH_-(\psi_n^{(-)}) = E_n^-(A\psi_n^{(-)}). \end{aligned} \quad (3)$$

Hence, except for the ground state that obeys $A\psi_0^{(-)} = 0$, for all other states $\psi_n^{(-)}$ of H_- there exists a state $\psi_{n-1}^{(+)} \propto A\psi_n^{(-)}$ of H_+ with exactly the same energy, i.e., $E_n^+ = E_{n-1}^-$, where $n = 0, 1, 2, \dots$. Conversely, one also has $A^\dagger\psi_n^{(+)} \propto \psi_{n+1}^{(-)}$.

Thus, if the eigenvalues and the eigenfunctions of H_- were known, one would automatically obtain the eigenvalues and the eigenfunctions of H_+ , which is in general a completely different Hamiltonian.

Now, let us consider the special case where $V_-(x, a_0)$ is a shape-invariant potential. For such systems, potentials $V_+(x, a_0) = V_-(x, a_1) + R(a_0)$. Their superpotential W obeys the integrability condition of Eq. (1). Since potentials $V_+(x, a_0)$ and $V_-(x, a_1)$ differ only in an additive constant, their common ground-state wave function is given by

$\psi_0^{(-)}(x, a_1) \sim \exp[-\int_{x_0}^x W(x, a_1)dx]$. [The ground-state energy of $H_+(x, a_0)$ is $R(a_0)$, because the ground-state energy of $V_-(x, a_1)$ vanishes.] Now using SUSY-QM algebra, the first excited state of $H_-(x, a_0)$ is given by $A^\dagger(x, a_0)\psi_0^{(-)}(x, a_1)$ and the corresponding eigenvalue is $R(a_0)$. By iterating this procedure, the $(n+1)$ th excited state is given by

$$\psi_{n+1}^{(-)}(x, a_0) \sim A^\dagger(a_0)A^\dagger(a_1)\cdots A^\dagger(a_n)\psi_0^{(-)}(x, a_n),$$

and corresponding eigenvalues are given by

$$E_0^{(-)} = 0; \quad \text{and} \quad E_n^{(-)} = \sum_{k=0}^{n-1} R(a_k) \quad \text{for } n > 0.$$

[To avoid notational complexity, we have suppressed the x dependence of operators $A(x, a_0)$ and $A^\dagger(x, a_0)$.] Thus, for a shape-invariant potential, one can obtain the entire spectrum of H_- by the algebraic methods of SUSY-QM.

Most of the known exactly solvable problems possess a spectrum generating algebra (SGA) [2,3,13]. We would like to find out if there is any connection between SGA and shape invariance of these systems. As we shall see later, the type of SGA that is most relevant to us is known as potential algebra, studied extensively by Alhassid and co-workers [2,3]. In potential algebra, the Hamiltonian of the system is written in terms of the Casimir operator (C_2) of the algebra, and the energy of states specified by an eigenvalue ω of C_2 is fixed. Different states with fixed ω represent eigenstates of a set of Hamiltonians that differ only in values of parameters and share a common energy. For a system with an $so(2,1)$ potential algebra, the different values of parameters are eigenvalues of operator J_3 , chosen to form a complete set of commuting observables. This is very similar to the case of shape-invariant potentials. In the next section, we will attempt to establish this connection in a more concrete fashion. In fact, for a set of solvable quantum-mechanical systems we shall explicitly show that shape invariance leads to a potential algebra.

III. SHAPE INVARIANCE AND CONNECTION TO ALGEBRA

Let us consider a generic shape-invariant potential $V_-(x, a_0)$ with a translational change of parameters $a_{m+1} = a_m + \delta = a_0 + (m+1)\delta$, where δ is a constant. For the superpotential $W(x, a_m) \equiv W(x, m)$, the shape-invariance condition is

$$W^2(x, m) + W'(x, m) = W^2(x, m+1) - W'(x, m+1) + R(m). \quad (4)$$

It is natural to ask whether the change of parameters can be formally accomplished by the action of a ladder-type operator. With this in mind, we define an operator $J_3 = -i\partial/\partial\phi \equiv -i\partial_\phi$, analogous to the z component of the angular-momentum operator. It acts upon functions in the space described by two coordinates x and ϕ , and its eigenvalues m will play the role of the parameter of the potential. We also define two more operators, J^- and its Hermitian conjugate J^+ by

$$J^\pm = e^{\pm i\phi} \left[\pm \frac{\partial}{\partial x} - W(x, -i\partial_\phi \pm \frac{1}{2}) \right]. \quad (5)$$

The factors $e^{\pm i\phi}$ in J^\pm ensure that they indeed behave as ladder operators for the quantum number m . (The factor $\pm \frac{1}{2}$ merely ‘‘symmetrizes’’ the J^\pm ’s.) Operators J^\pm are basically of the same form as the A^\pm operators of SUSY-QM, except that the parameter m of the superpotential W is replaced by operators $(J_3 \pm \frac{1}{2})$. Explicit computation shows that

$$[J_3, J^\pm] = \pm J^\pm, \quad (6)$$

and hence operators J^\pm change the eigenvalues of the J_3 operator by unity, similar to the ladder operators of angular momentum [so(3)]. Now let us determine the commutator $[J^+, J^-]$:

$$\begin{aligned} [J^+, J^-] &= J^+ J^- - J^- J^+ \\ &= \left[-\frac{\partial^2}{\partial x^2} + W^2(x, J_3 - \frac{1}{2}) - W'(x, J_3 - \frac{1}{2}) \right] \\ &\quad - \left[-\frac{\partial^2}{\partial x^2} + W^2(x, J_3 + \frac{1}{2}) + W'(x, J_3 + \frac{1}{2}) \right] \\ &= -R(J_3 + \frac{1}{2}), \end{aligned} \quad (7)$$

where we have used the shape-invariance condition (4). Thus, we see that shape invariance enables us to close the algebra of J_3 and J^\pm to

$$[J_3, J^\pm] = \pm J^\pm, \quad [J^+, J^-] = -R(J_3 + \frac{1}{2}). \quad (8)$$

Now, if the function $R(J_3)$ is linear in J_3 , the algebra of Eq. (8) reduces to that of so(3) or so(2,1) [9]. Several SIP’s are of this type, among them are the Morse, Scarf I, Scarf II, and generalized Pöschl-Teller potentials. For these potentials, $R(J_3 + \frac{1}{2}) = 2J_3$ [4], and Eq. (8) reduces to an so(2,1) algebra and hence establishes a connection between shape invariance and potential algebra. With a slightly different formalism, Balantekin arrived at a similar conclusion for these SIP’s [14]. However, there are many other important systems like Coulomb, Eckart, etc. where $R(a_0)$ is not linear in a_0 , and these cases will be discussed later.

IV. DIFFERENTIAL REALIZATION OF so(2,2)

Before establishing a connection between a general Nanzon Hamiltonian and an so(2,2) potential algebra, we will discuss a realization of so(2,2) algebra in terms of differential operators on a (2,2)-hyperboloid. For consistency, we use the formalism and the notations of Refs. [2,3].

A (2,2)-hyperboloid is defined by

$$\begin{aligned} x_1 &= \rho \cosh \chi \cos \phi, & x_2 &= \rho \cosh \chi \sin \phi, \\ x_3 &= \rho \sinh \chi \cos \theta, & x_4 &= \rho \sinh \chi \sin \theta, \end{aligned} \quad (9)$$

where ϕ and θ are rotation angles in the x_1, x_2 and x_3, x_4 planes, respectively [$0 \leq \phi, \theta < 2\pi$]. Six generators of the algebra, J_i and K_i ($i = 1, \dots, 3$) can be chosen as

$$\begin{aligned} J_1 &= (x_2 p_3 + x_3 p_2), & J_2 &= -(x_1 p_3 + x_3 p_1), \\ J_3 &= (x_1 p_2 - x_2 p_1), \\ K_1 &= (x_1 p_4 + x_4 p_1), & K_2 &= (x_2 p_4 + x_4 p_2), \\ K_3 &= (x_3 p_4 - x_4 p_3). \end{aligned} \quad (10)$$

Operators p_i represent derivatives $-i(\partial/\partial x_i)$. The algebraic relations obeyed by these operators are given by

$$\begin{aligned} [J_1, J_2] &= -iJ_3, & [J_2, J_3] &= iJ_1, & [J_3, J_1] &= iJ_2, \\ [K_1, K_2] &= -iJ_3, & [K_2, K_3] &= iJ_1, & [K_3, K_1] &= iJ_2, \\ [J_1, K_2] &= -iK_3, & [J_2, K_3] &= iK_1, & [J_3, K_1] &= iK_2, \\ [K_1, J_2] &= -iK_3, & [K_2, J_3] &= iK_1, & [K_3, J_1] &= iK_2. \end{aligned} \quad (11)$$

The above algebra can be decomposed in terms of two commuting so(2,1) algebras generated by

$$A_i = \frac{1}{2}(J_i + K_i), \quad B_i = \frac{1}{2}(J_i - K_i). \quad (12)$$

These operators commute; i.e., $[A_i, B_j] = 0$. Using Eqs. (9) and (11), the differential realization can be written explicitly as [2,3]

$$\begin{aligned} A^\pm &\equiv A_1 \pm iA_2 = \frac{1}{2} e^{\pm i(\phi + \theta)} \\ &\quad \times \left[\mp \frac{\partial}{\partial \chi} + \tanh \chi(-i\partial_\phi) + \coth \chi(-i\partial_\theta) \right], \\ A_3 &= -\frac{i}{2}(\partial_\phi + \partial_\theta), \\ B^\pm &\equiv B_1 \pm iB_2 = \frac{1}{2} e^{\pm i(\phi - \theta)} \\ &\quad \times \left[\mp \frac{\partial}{\partial \chi} + \tanh \chi(-i\partial_\phi) + \coth \chi(+i\partial_\theta) \right], \\ B_3 &= -\frac{i}{2}(\partial_\phi - \partial_\theta). \end{aligned} \quad (13)$$

The so(2,1) algebra obeyed by these operators is

$$[A_3, A^\pm] = \pm A^\pm, \quad [A^+, A^-] = -2A_3$$

and a similar one for the B ’s. The Casimir operator C_2 is given by

$$\begin{aligned} C_2 &= 2(A_3^2 - A_+ A_- - A_3) + 2(B_3^2 - B_+ B_- - B_3) \\ &= \left[\frac{\partial^2}{\partial \chi^2} + (\tanh \chi + \coth \chi) \frac{\partial}{\partial \chi} + \operatorname{sech}^2 \chi(-i\partial_\phi)^2 \right. \\ &\quad \left. - \operatorname{cosech}^2 \chi(-i\partial_\theta)^2 \right]. \end{aligned} \quad (14)$$

Operators A_3, B_3 , and C_2 can be simultaneously diagonalized, and their actions on their common eigenstates are given by

$$\begin{aligned}
 C_2|\omega, m_1, m_2\rangle &= \omega(\omega + 2)|\omega, m_1, m_2\rangle, \\
 A_3|\omega, m_1, m_2\rangle &= m_1|\omega, m_1, m_2\rangle, \\
 B_3|\omega, m_1, m_2\rangle &= m_2|\omega, m_1, m_2\rangle.
 \end{aligned}
 \tag{15}$$

It is worth mentioning at this point that the Casimir operator given above is indeed self-adjoint with respect to a measure $\sinh \chi \cosh \chi d\chi d\phi d\theta$.

V. NATANZON POTENTIALS

The Schrödinger equation for any Natanzon potential can be reduced by a point canonical transformation (a general similarity transformation followed by an appropriate change of independent variable) [10,15,16] to the hypergeometric equation. A general potential $U(r)$ of the Natanzon type is implicitly defined by [1]

$$U[z(r)] = \frac{-fz(1-z) + h_0(1-z) + h_1z}{R(z)} - \frac{1}{2}\{z, r\}, \tag{16}$$

where $R(z) = az^2 + b_0z + c_0 = a(1-z)^2 - b_1(1-z) + c_1$ and $f, h_0, h_1, a, b_0, b_1, c_0, c_1$ are constants. The Schwarzian derivative $\{z, r\}$ is defined by

$$\{z, r\} \equiv \frac{d^3z/dr^3}{dz/dr} - \frac{3}{2} \left[\frac{d^2z/dr^2}{dz/dr} \right]^2. \tag{17}$$

The relationship between variables z ($0 < z < 1$) and r is implicitly given by

$$\left(\frac{dz}{dr} \right) = \frac{2z(1-z)}{\sqrt{R(z)}}. \tag{18}$$

To avoid a singularity in $U(z(r))$, one assumes that $R(z)$ has no singularity in the domain $(0,1)$. The Schrödinger equation is given by

$$\left[\frac{d^2}{dr^2} + \left\{ \left(\frac{dz}{dr} \right)^2 I(z) + \frac{1}{2}\{z, r\} \right\} \right] = 0,$$

where

$$I(z) = \frac{(1-\lambda_0^2)(1-z) + (1-\lambda_1^2)z + (\mu^2-1)z(1-z)}{4z^2(1-z)^2}$$

and

$$\begin{aligned}
 (1-\mu^2) &= aE - f, & (1-\lambda_0^2) &= c_0E - h_0, \\
 (1-\lambda_1^2) &= c_1E - h_1.
 \end{aligned}
 \tag{19}$$

To connect the Casimir operator C_2 of the $so(2,2)$ algebra [Eq. (14)] to the general Natanzon potential, we will first perform a similarity transformation on C_2 by a function F and then follow that up by an appropriate change of variable $\chi \rightarrow g(r)$. Under the similarity transformation,

$$\begin{aligned}
 \frac{d}{d\chi} &\rightarrow F \frac{d}{d\chi} F^{-1} = \left(\frac{d}{d\chi} - \frac{\dot{F}}{F} \right), \\
 \frac{d^2}{d\chi^2} &\rightarrow \left(\frac{d^2}{d\chi^2} - \frac{2\dot{F}}{F} \frac{d}{d\chi} + \frac{2\dot{F}^2}{F^2} - \frac{\ddot{F}}{F} \right),
 \end{aligned}$$

where dots represent derivatives with respect to χ . The Casimir operator C_2 of Eq. (14) transforms as

$$\begin{aligned}
 C_2 \rightarrow \tilde{C}_2 &= \left[\frac{d^2}{d\chi^2} + \left(\tanh \chi + \coth \chi - \frac{2\dot{F}}{F} \right) \frac{d}{d\chi} \right. \\
 &+ \frac{2\dot{F}^2}{F^2} - \frac{\ddot{F}}{F} - (\tanh \chi + \coth \chi) \frac{\dot{F}}{F} \\
 &\left. + \operatorname{sech}^2 \chi (-i\partial_\phi)^2 - \operatorname{cosech}^2 \chi (-i\partial_\theta)^2 \right].
 \end{aligned}
 \tag{20}$$

Now, let us carry out a change of variable from χ to r via $\chi = g(r)$. We are going to denote differentiation with respect to r by a prime. Operators $d/d\chi$ and $d^2/d\chi^2$ transform as

$$\frac{d}{d\chi} = \frac{1}{g'} \frac{d}{dr}, \quad \frac{d^2}{d\chi^2} = \frac{1}{g'^2} \left[\frac{d^2}{dr^2} - \frac{g''}{g'} \frac{d}{dr} \right].$$

The operator \tilde{C}_2 now transforms into

$$\begin{aligned}
 \tilde{C}_2 &= \frac{1}{g'^2} \left[\frac{d^2}{dr^2} + \left\{ -\frac{g''}{g'} - \frac{2F''}{F} + g'(\tanh g + \coth g) \right\} \right. \\
 &\times \frac{d}{dr} + \frac{2F'^2}{F^2} - \frac{F''}{F} + \frac{F'g''}{Fg'} - \frac{F'g'}{F}(\tanh g + \coth g) \\
 &\left. + g'^2(\operatorname{sech}^2 g (-i\partial_\phi)^2 - \operatorname{cosech}^2 g (-i\partial_\theta)^2) \right].
 \end{aligned}
 \tag{21}$$

In order for $g'^2 \tilde{C}_2$ to be a Schrödinger Hamiltonian, we require the expression inside the curly brackets in Eq. (21) to vanish. This constrains the relationship between the two functions F and g to be

$$-\frac{g''}{g'} - \frac{2F'}{F} + g'(\tanh g + \coth g) = 0, \tag{22}$$

which yields

$$F \sim \left(\frac{\sinh(2g)}{g'} \right)^{1/2}. \tag{23}$$

Thus the operator \tilde{C}_2 transforms into

$$\tilde{C}_2 = \frac{1}{g'^2} \left[\frac{d^2}{dr^2} + g'^2 \left(\frac{(1 - \tanh^2 g)^2 - 4 \tanh^2 g}{4 \tanh^2 g} \right) + \frac{1}{2} \{g, r\} + g'^2 (\operatorname{sech}^2 g (-i \partial_\phi)^2 - \operatorname{cosech}^2 g (-i \partial_\theta)^2) \right]. \tag{24}$$

This Casimir operator now has a form of

$$\tilde{C}_2 = -\frac{1}{g'^2} H,$$

where H is a one-dimensional Hamiltonian with the potential $U(r)$ given by

$$U(z(r)) = \frac{ER + \left[-\frac{7}{4} + \frac{5}{2}z - \frac{7}{4}z^2 \right] - z(1-z)(-i \partial_\phi)^2 + (1-z)(-i \partial_\theta)^2}{R} - \frac{1}{2} \{z, r\} = \{ -[aE - \frac{7}{4} + (-i \partial_\phi)^2]z(1-z) + [c_0E - \frac{7}{4} + (-i \partial_\theta)^2](1-z) + [(a + b_0 + c_0)E - 1] \} / R(z) - \frac{1}{2} \{z, r\}. \tag{25}$$

We have used

$$g' = \frac{dg}{dz} z' = \frac{1}{2\sqrt{z(1-z)}} \frac{2z(1-z)}{R} = \sqrt{\frac{z}{R}}.$$

Now, with the following identification:

$$\begin{aligned} f &= aE - \frac{7}{4} + (-i \partial_\phi)^2, \\ h_0 &= c_0E - \frac{7}{4} + (-i \partial_\theta)^2, \\ h_1 &= (a + b_0 + c_0)E - 1, \end{aligned} \tag{26}$$

$$E - U(r) = g'^2 \left(\frac{(1 - \tanh^2 g)^2 - 4 \tanh^2 g}{4 \tanh^2 g} \right) + \frac{1}{2} \{g, r\} + g'^2 [\operatorname{sech}^2 g (-i \partial_\phi)^2 - \operatorname{cosech}^2 g (-i \partial_\theta)^2].$$

Now, for this potential to take the form of a general Natanzon potential, we have to relate variables g and z in such a way that the potential in terms of z is given by Eq. (16). Since the potential has to be a ratio of two quadratic functions of z , this is accomplished with the identification $z = \tanh^2 g$, which leads to

the potential of Eq. (25) indeed has the form of a general Natanzon potential [Eq. (16)].

VI. SHAPE-INVARIANT NATANZON POTENTIALS FROM POTENTIAL ALGEBRA

At this point we go back to the operators A^\pm [Eq. (13)] and see how they transform under the similarity transformation given by $F \sim \{[\sinh(2g)]/g'\}^{1/2} \sim \sqrt{z/z'}$. This transformation carries operators A^\pm to

TABLE I. All allowed values of α, β and the superpotentials that they generate. Constants \tilde{m}_i ($i=1,2$) are linear combinations of m_1 and m_2 of Eq. (15). Simple harmonic-oscillator, Morse, and Coulomb potentials are not included in this list, as they can be considered to be limiting cases of potentials presented here [10].

α	β	$z(r)$	Superpotential	Potential
0	0	$z = e^{-r}$	$\tilde{m}_1 \coth \frac{r}{2} + \tilde{m}_2$	Eckart
0	$-\frac{1}{2}$	$z = \sin^2 \frac{r}{2}$	$\tilde{m}_1 \operatorname{cosec} r + \tilde{m}_2 \cot r$	Gen. Pöschl-Teller trigonometric
0	-1	$z = 1 - e^{-r}$	$\tilde{m}_1 \coth \frac{r}{2} + \tilde{m}_2$	Eckart
$-\frac{1}{2}$	0	$z = \operatorname{sech}^2 \frac{r}{2}$	$\tilde{m}_1 \operatorname{cosech} r + \tilde{m}_2 \coth r$	Pöschl-Teller II
$-\frac{1}{2}$	$-\frac{1}{2}$	$z = \tanh^2 \frac{r}{2}$	$\tilde{m}_1 \tanh \frac{r}{2} + \tilde{m}_2 \coth \frac{r}{2}$	Gen. Pöschl-Teller
-1	0	$z = 1 + \tanh \frac{r}{2}$	$\tilde{m}_1 \tanh \frac{r}{2} + \tilde{m}_2$	Rosen Morse

$$A^\pm \rightarrow \tilde{A}^\pm = \frac{e^{\pm i(\phi+\theta)}}{2} \left[\mp \left(\frac{d}{d\chi} + \frac{1}{2z'} \frac{dz'}{d\chi} - \frac{1}{2z} \frac{dz}{d\chi} \right) + \tanh \chi(-i\partial_\phi) + \coth \chi(-i\partial_\theta) \right]. \quad (27)$$

If the expression $[(1/2z')(dz'/d\chi) - (1/2z)(dz/d\chi)]$ can be written as a linear combination of $\tanh \chi$ and $\coth \chi$, operators \tilde{A}^\pm can be cast in a form similar to the operators J^\pm of Eq. (5), and the connection with shape invariance is established.

Hence to get shape-invariant potentials we require

$$\left(\frac{1}{2z'} \frac{dz'}{d\chi} - \frac{1}{2z} \frac{dz}{d\chi} \right) = \alpha \tanh \chi + \beta \coth \chi.$$

This leads to $z' = z^{1+\beta}(1-z)^{-\alpha-\beta}$, which is another restriction on the relationship between variables z and r . Since these variables are already constrained by Eq. (18), only a handful of solutions would be compatible with both restrictions. Thus $z(r)$'s that are compatible with both equations are given by

$$z^{1+\beta}(1-z)^{-\alpha-\beta} = \frac{2z(1-z)}{\sqrt{R(z)}}, \quad (28)$$

where $R(z)$ is a quadratic function of z . After some computation, we find that there are only a finite number of values of α, β that satisfy Eq. (28). These values are listed in Table I, and they exhaust all known shape-invariant potentials that lead to the hypergeometric equation. It is also interesting to

note that while the potential algebra of a general Natanzon system is $so(2,2)$, and requires two sets of raising and lowering operators A^\pm and B^\pm , all known shape-invariant potentials need only one such set. For all SIP's of Table 4.1 of Ref. [4], one finds that all partner potentials are connected by a change of just one independent parameter (although other parameters which do not change are also present). Thus there is a series of potentials that only differ in one parameter. From the potential algebra perspective, all these potentials differ only by the eigenvalue of an operator that is a linear combination of A_3 and B_3 , and all are characterized by a common eigenvalue of C_2 . Thus, these shape-invariant potentials can be associated with an $so(2,1)$ potential algebra generated by operators A^+, A^-, B^+ and the same linear combination of A_3 and B_3 .

Note added in the proof. Recently, we became aware of work done by Cordero and Salamo [17]. In these papers, the authors show that the Natanzon potential has a spectrum-generating algebra given by $so(2,1)$. However, for shape-invariant systems, the potential algebra [2,3] generated by operators with a first-order derivative is more relevant. The Casimir operator of the potential algebra is essentially the same as the Hamiltonian of the shape-invariant model.

ACKNOWLEDGMENTS

A.G. acknowledges a research leave from Loyola University, Chicago, which made his involvement in this work possible. We would like to thank Professor Y. Alhassid for clarifying several points. One of us (A.G.) would also like to thank the Physics Department of the University of Illinois for their warm hospitality. Partial financial support from the U.S. Department of Energy is gratefully acknowledged.

-
- [1] G. A. Natanzon, *Theor. Math. Phys.* **38**, 146 (1979) [*Teor. Math. Fiz.* **38**, 219 (1979)].
- [2] Y. Alhassid, F. Gürsey, and F. Iachello, *Ann. Phys. (Leipzig)* **148**, 346 (1983).
- [3] J. Wu and Y. Alhassid, *J. Math. Phys.* **31**, 557 (1990).
- [4] F. Cooper, A. Khare, and U. Sukhatme, *Phys. Rep.* **251**, 268 (1995), and references therein.
- [5] L. Infeld and T. E. Hull, *Rev. Mod. Phys.* **23**, 21 (1951).
- [6] L. E. Gendenshtein, *Pisma Zh. Eksp. Teor. Fiz.* **38**, 299 (1983) [*JETP Lett.* **38**, 356 (1983)].
- [7] V. P. Spiridonov, *Phys. Rev. Lett.* **69**, 398 (1992).
- [8] D. Barclay, R. Dutt, A. Gangopadhyaya, A. Khare, A. Pagnamenta, and U. Sukhatme, *Phys. Rev. A* **48**, 2786 (1993).
- [9] A. Gangopadhyaya, J. V. Mallow, and U. P. Sukhatme, in *Proceedings of Workshop on Supersymmetric Quantum Mechanics and Integrable Models, 1997*, edited by Henrik Aratyn *et al.* (Springer-Verlag, Berlin, 1998).
- [10] A. Gangopadhyaya, P. Panigrahi, and U. Sukhatme, *Helv. Phys. Acta* **67**, 363 (1994).
- [11] A. Comtet, A. Bandrauk, and D. K. Campbell, *Phys. Lett. B* **150**, 381 (1985).
- [12] J. S. Dowker, *J. Phys. A* **3**, 451 (1970); *Ann. Phys. (Leipzig)* **62**, 561 (1971); K. Funahashi, T. Kashiwa, S. Sakoda, and K. Fujii, *J. Math. Phys.* **36**, 3232 (1995); **36**, 4590 (1995).
- [13] A. O. Barut, A. Inomata, and R. Wilson, *J. Phys. A* **20**, 4075 (1987); **20**, 4083 (1987); M. J. Englefield, *J. Math. Phys.* **28**, 827 (1987); M. J. Englefield and C. Quesne, *J. Phys. A* **24**, 3557 (1987); R. D. Tangerman and J. A. Tjon, *Phys. Rev. A* **48**, 1089 (1993).
- [14] A. B. Balantekin, *Phys. Rev. A* **57**, 4188 (1998).
- [15] F. Cooper, J. N. Ginnocchio, and A. Wipf, *J. Phys. A* **22**, 3707 (1989).
- [16] R. De, R. Dutt, and U. Sukhatme, *J. Phys. A* **25**, L843 (1992).
- [17] P. Cordero and S. Salamo, *Found. Phys.* **23**, 675 (1993); *J. Math. Phys.* **35**, 3301 (1994).