



2011

Exceptional Orthogonal Polynomials, QHJ Formalism and SWKB Quantization Condition

S Ree Ranjani

P K. Panigrahi

A Khare

A K. Kapoor

Asim Gangopadhyaya

Loyola University Chicago, agangop@luc.edu

Author Manuscript

This is a pre-publication author manuscript of the final, published article.

Recommended Citation

Ranjani, SR, PK Panigrahi, A Khare, AK Kapoor, and A Gangopadhyaya. "Exceptional Orthogonal Polynomials, QHJ Formalism and SWKB Quantization Condition." *Journal of Physics A: Mathematical and Theoretical* 45, 2012.

This Article is brought to you for free and open access by the Faculty Publications at Loyola eCommons. It has been accepted for inclusion in Physics: Faculty Publications and Other Works by an authorized administrator of Loyola eCommons. For more information, please contact ecommons@luc.edu.



This work is licensed under a [Creative Commons Attribution-Noncommercial-No Derivative Works 3.0 License](https://creativecommons.org/licenses/by-nc-nd/3.0/).

© IOP Publishing, 2011.

Exceptional orthogonal polynomials, QHJ formalism and SWKB quantization condition

S. Sree Ranjani^{a 1}, P. K. Panigrahi^{b 2}, A. Khare^{c 3}, A. K. Kapoor^{a 4} and A. Gangopadhyaya^{d 5}

^aSchool of Physics, University of Hyderabad, Hyderabad, 500 046, India.

^bIndian Institute of Science Education and Research (IISER) Kolkata, Salt Lake, Kolkata, 700 106, India.

^cIndian Institute of Science Education and Research (IISER) Pune, 411021, India.

^dLoyola University Chicago, Department of Physics, Chicago, IL 60660.

Abstract

We study the quantum Hamilton-Jacobi (QHJ) equation of the recently obtained exactly solvable models, related to the newly discovered exceptional polynomials and show that the QHJ formalism reproduces the exact eigenvalues and the eigenfunctions. The fact that the eigenfunctions have zeros and poles in complex locations leads to an unconventional singularity structure of the quantum momentum function $p(x)$, the logarithmic derivative of the wave function, which forms the crux of the QHJ approach to quantization. A comparison of the singularity structure for these systems with the known exactly solvable and quasi-exactly solvable models reveals interesting differences. We find that the singularity structure of the momentum function for these new potentials lies between the above two distinct models, sharing similarities with both of them. This prompted us to examine the exactness of the supersymmetric WKB (SWKB) quantization condition. The interesting singularity structure of $p(x)$ and of the superpotential for these models has important consequences for the SWKB rule and in our proof of its exactness for these quantal systems.

1 Introduction

New infinite sets of solvable quantum mechanical potentials have been recently constructed by Odake and Sasaki [1],[2], using the Darboux-Crum transformations [3],[4]. They deformed the standard radial oscillator and Darboux-Pöschl-Teller potentials using the Laguerre and the Jacobi polynomial eigenfunctions of degree ℓ . The solutions of the new potentials are in terms of the newly discovered Laguerre- and the Jacobi- X_ℓ type exceptional orthogonal polynomials respectively, which have been recently introduced by Gómez-Ullate *et.al.*, [5],[6]. The characteristic feature of this new class of polynomials is that the series starts with a polynomial of degree ℓ ($\ell = 1, 2, \dots$) and still can form a complete set with respect to a positive-definite measure. This is unlike the classical orthogonal polynomial series, which require a constant to form a complete set. In addition to this feature, the eigenfunctions of these new potentials have singularities at complex locations, associated with the zeros of the eigenpolynomials of the standard potentials

¹s.sreeranjani@gmail.com

²prasanta@iiserkol.ac.in

³khare@iiserpune.ac.in

⁴akksp@uohyd.ernet.in

⁵A. Gangopadhyaya

from which they are derived. These standard polynomials seem to play a crucial role in the construction of these new potentials. The fact that the new Laguerre- and the Jacobi- X_ℓ type polynomials occur as solutions to the Sturm-Liouville problem with rational coefficients, associated to the standard Laguerre and Jacobi polynomials, is very intriguing. In this light the quantum Hamilton-Jacobi (QHJ) analysis [7], [8], which uses the singularity structure of the logarithmic derivative of the wave function to obtain the eigenvalues and eigenfunctions for a given potential, can reveal interesting features pertaining to these models.

In this paper, we analyze the set of potentials obtained by deforming the radial oscillator using the quantum Hamilton-Jacobi (QHJ) formalism and investigate the singularity structure of the logarithmic derivative of the wave function, $\psi(x)$:

$$p(x) = -i\hbar \frac{\partial_x \psi(x)}{\psi(x)}, \quad (1)$$

known as the quantum momentum function (QMF). The knowledge of the singularity structure of the QMF, coupled with the exact quantization condition,

$$\frac{1}{2\pi} \oint_C p(x) dx = n\hbar, \quad (2)$$

defined within this formalism, allows us to arrive at the required solutions [9]-[12]. Here, n gives the number of nodes of $\psi(x)$ and the contour C encloses these nodes located in the classical region, in between the two turning points x_1 and x_2 in the complex x -plane. In the present study, in addition to obtaining the energy eigenvalues and eigenfunctions, we concentrate on bringing out the new features of the singularity structure of the QMF of these potentials. This, when compared to that of the models studied earlier, namely the exactly solvable (ES) models, whose solutions are in terms of the classical polynomials, and the quasi-exactly solvable (QES) models revealed interesting differences. This led us to a careful analysis of the exactness or otherwise of the supersymmetric WKB (SWKB) quantization condition [13], [14] for these new models, because it was shown in [15] that the singularity structure of the QMF provided a link to the exactness of the SWKB integral. Here, Bhalla et.al., showed that the SWKB condition is exact for all models, where the locations and the residues of the poles in the non-classical regions of the SWKB integrand and the QMF matched identically. Hence, it will be interesting to see if a similar analysis works for these new models, especially when their singularity structure differs from that of the conventional models.

In the following section, we give a brief account of the QHJ formalism [7]-[12], followed by the analysis of the generalized deformed radial oscillator potential within this method. We then compare the singularity structures of various models, in order to identify the similarities or differences, between the ES and QES models and the present ones. In section 3 we investigate the exactness of the SWKB condition for the deformed radial oscillators and subsequently conclude in the last section, with directions for further research.

2 QHJ formalism and the deformed radial oscillators

In the QHJ formalism, x is treated as a complex variable and we use the Riccati equation for $q (= ip(x))$ ($\hbar = 2m = 1$)

$$q^2 + \partial_x q + (E - V(x)) = 0, \quad (3)$$

to obtain the required solutions. Using (1) and substituting $q = \frac{\psi'(x)}{\psi(x)}$ in the above equation gives us the Schrödinger equation

$$-\frac{d^2\psi(x)}{dx^2} + V(x)\psi(x) = E\psi(x), \quad (4)$$

which establishes the connection between QHJ approach and the conventional ones. In this approach, the singularities of the momentum function q play the key role in obtaining the spectra, without the need to solve for the eigenfunctions.

The singularities of q are of two types namely, the fixed and the moving singularities. The fixed singularities correspond to the singularities of the potential and their location is independent of the energy and can be located by inspection from (3). The moving singularities are poles, which correspond to the zeros of the wave function $\psi(x)$. It is well known that the location of the nodes of the wave function changes with energy. Therefore, the location of these poles depends on the initial condition and hence they are named moving poles. The number and the location of these poles cannot be inferred from the differential equation by inspection. These poles turn out to be finite in number for all the ES models studied [10], [12]. This is equivalent to the fact that the point at infinity, at best, is an isolated singularity. The location of all these singularities and their residues are used to evaluate the integral in (2), which gives the expression for the eigenvalues for the ES models [10] and the quasi-exact solvability condition for the QES models [11]. In addition, the knowledge of the singularity structure of q allows us to obtain the form of QMF and hence the eigenfunctions using (1). For more details, we refer the reader to our earlier papers [9]-[12] and we proceed to analyze the potentials under study.

Deformed radial oscillator potentials

Here, we present a brief account of the set containing infinite number of new shape invariant potentials (SIPs) $V_\ell(x)$, indexed by ℓ ($\ell = 1, 2, \dots$) described in [1], [2]. The potentials $V_\ell(x)$ constitute the hierarchy of supersymmetric potentials and are translationally shape invariant [13], [16]. Moreover, within each set, putting $\ell = 0$ gives the respective standard potential and $\ell = 1$ corresponds to the potentials constructed by Quesne [17]. In [18], all the SIPs including the newly deformed potentials have been obtained from a general analysis of shape invariance requirement. Here, we analyze the set of potentials obtained by deforming the standard radial oscillator. For every potential in this set, indexed by ℓ , a prepotential

$$\omega_\ell(x; g) = \omega_0(x; g + \ell) + \ln \frac{\xi_\ell(x^2; g + 1)}{\xi_\ell(x^2; g)}, \quad g > 0 \quad (5)$$

is defined, where

$$\omega_0(x; g) = -\frac{1}{2}x^2 + g \log x, \quad 0 < x < \infty, \quad (6)$$

$$\xi_\ell(x^2; g) = L_\ell^{(g+\ell-\frac{3}{2})}(-x^2). \quad (7)$$

Here $L_\ell^{(g+\ell-\frac{3}{2})}(x)$ is the associated Laguerre polynomial. The corresponding superpotential is given by

$$W_\ell(x) = -\partial_x w_\ell(x; g) = x - \frac{g+\ell}{x} - \frac{\partial_x \xi_\ell(x^2; g+1)}{\xi_\ell(x^2; g+1)} + \frac{\partial_x \xi_\ell(x^2; g)}{\xi_\ell(x^2; g)} \quad (8)$$

and the corresponding potential, $V_\ell(x)$, with zero ground state energy is given by

$$V_\ell(x) = (\partial_x w_\ell(x; g))^2 + \partial_{xx} w_\ell(x; g). \quad (9)$$

Substituting $\omega_\ell(x; g)$ in the above equation and simplifying we obtain,

$$\begin{aligned} V_\ell(x) &= U_0^2(x) + \partial_x U_0(x) + \frac{\partial_{xx} \xi_\ell(x^2; g+1)}{\xi_\ell(x^2; g+1)} \\ &+ 2 \left(-x + \frac{g+\ell}{x} - \frac{\partial_x \xi_\ell(x^2; g)}{\xi_\ell(x^2; g)} \right) \frac{\partial_x \xi_\ell(x^2; g+1)}{\xi_\ell(x^2; g+1)}, \end{aligned} \quad (10)$$

where

$$U_0(x) = -x + \frac{g+\ell}{x} - \frac{\partial_x \xi_\ell(x^2; g)}{\xi_\ell(x^2; g)}. \quad (11)$$

From the differential equation and the recurrence relations of the associated Laguerre polynomials [19],[2], we obtain

$$\frac{\partial_{xx} \xi_\ell(x^2; g+1)}{\xi_\ell(x^2; g+1)} = -2 \left(-x + \frac{g+\ell}{x} \right) \frac{\partial_x \xi_\ell(x^2; g+1)}{\xi_\ell(x^2; g+1)} + 4\ell \quad (12)$$

and

$$\left(\frac{\partial_x \xi_\ell(x^2; g)}{\xi_\ell(x^2; g)} \right) \left(\frac{\partial_x \xi_\ell(x^2; g+1)}{\xi_\ell(x^2; g+1)} \right) = 2x \frac{\partial_x \xi_\ell(x^2; g+1)}{\xi_\ell(x^2; g)} \left(1 - \frac{\partial_x \xi_\ell(x^2; g)}{\xi_\ell(x^2; g+1)} \right) \quad (13)$$

respectively. The use of these equations in (10), reduces the potential to

$$V_\ell(x) = U_0^2(x) + \partial_x U_0(x) - 4x \frac{\partial_x \xi_\ell(x^2; g+1)}{\xi_\ell(x^2; g)} + 4\ell. \quad (14)$$

Substituting (14) in (3) gives us the QHJ equation

$$q_\ell^2 + \partial_x q_\ell + E = U_0^2(x) + \partial_x U_0(x) - 4x \frac{\partial_x \xi_\ell(x^2; g+1)}{\xi_\ell(x^2; g)} + 4\ell. \quad (15)$$

Here, q_ℓ corresponds to the momentum function associated to $V_\ell(x)$. From the above equation we can see that q_ℓ has a simple pole at the origin along with 2ℓ fixed poles corresponding to the zeros of $\xi_\ell(x^2; g)$. An important feature of all known ES models studied has been that the point at infinity is an isolated singularity [10]. The consequence of this is that the QMF has finite number of moving poles. For the new ES models under study we make the same assumption, *i.e.*, momentum function has a finite number, N , moving poles. From (15), we can see that q_ℓ behaves like x at infinity. This coupled

with the knowledge of its singularities and residues allows us to write q_ℓ in the form of a meromorphic function

$$q_\ell = -x + \frac{g + \ell}{x} - \sum_{i=1}^{2\ell} \frac{1}{x - a_i} + \sum_{j=1}^N \frac{1}{x - b_j} + \phi(x). \quad (16)$$

The first term on the right hand side of the above equation gives the large x behaviour of q_ℓ and the second term is the singular part arising from the pole at the origin. The numerator is the residue at the origin and is calculated by substituting the Laurent expansion of q_ℓ around the same :

$$q_\ell = \frac{b_1}{x} + a_0 + a_1x + \dots \quad (17)$$

in (15). Equating the coefficients of the same powers of x to zero, we obtain two values for b_1 , namely $\pm(g + \ell)$. The correct value is chosen using the boundary condition, in the limit $E \rightarrow 0$, $q_\ell \rightarrow -W_\ell(x)$. This gives the residue to be $(g + \ell)$. The third term in (16) gives the singular part associated with 2ℓ fixed poles, with residue -1 , corresponding to the zeros of the $\xi_\ell(x^2, g)$. It is easily seen that this is equivalent to $\frac{\partial_x \xi_\ell(x^2, g)}{\xi_\ell(x^2, g)}$. The fourth term corresponds to N simple moving poles with residue 1. Some of these are located on the real line and others off the real line. The ones which lie on the positive real line correspond to the nodes of the wave function. The contribution of the moving poles of q_ℓ is similarly written as $\frac{\partial_x P_N(x)}{P_N(x)}$, with $P_N(x)$ being a polynomial of degree N . The last term, $\phi(x)$, describes the analytic part of q_ℓ . From Liouville's theorem, $\phi(x)$ is a constant, which turns out to be zero from (15). Substituting (16) in (15) and after a few algebraic manipulations we get

$$\begin{aligned} \frac{\partial_{xx} P_N(x)}{P_N(x)} + 2 \left(-x + \frac{g + \ell}{x} - \frac{\partial_x \xi_\ell(x^2; g)}{\xi_\ell(x^2; g)} \right) \frac{\partial_x P_N(x)}{P_N(x)} \\ + 4x \frac{\partial_x \xi_\ell(x^2; g + 1)}{\xi_\ell(x^2; g)} - 4\ell + E = 0. \end{aligned} \quad (18)$$

For large x the leading term in the above equation is a constant, therefore equating it to zero, we obtain

$$E = 2N - 4\ell. \quad (19)$$

Substituting this in (18), we obtain

$$\begin{aligned} \partial_{xx} P_N(x) + 2 \left[-x + \frac{g + \ell}{x} - \frac{\partial_x \xi_\ell(x^2; g)}{\xi_\ell(x^2; g)} \right] \partial_x P_N(x) + \\ \left(4x \frac{\partial_x \xi_\ell(x^2; g + 1)}{\xi_\ell(x^2; g)} + 2N - 8\ell \right) P_N(x) = 0. \end{aligned} \quad (20)$$

The above equation can be written as

$$\frac{1}{\mathbf{w}_\ell^2} \frac{d}{dx} \left(\mathbf{w}_\ell^2(x) \frac{d}{dx} P_N(x) \right) + \left(-4x \frac{\partial_x \xi_\ell(x^2; g + 1)}{\xi_\ell(x^2; g)} + 2N - 8\ell \right) P_N(x) = 0 \quad (21)$$

where,

$$\begin{aligned}\mathbf{w}_\ell(x) &= \exp \left[\int \left(-x + \frac{g + \ell}{x} - \frac{\partial_x \xi_\ell(x^2; g)}{\xi_\ell(x^2; g)} \right) dx \right], \\ &= x^{(g+\ell)} \exp(-\frac{1}{2}x^2) / \xi_\ell(x^2; g).\end{aligned}\tag{22}$$

The differential equation (21) implies that \mathbf{w}_ℓ^2 is the weight function with respect to which the polynomials, $P_N(x)$, are orthogonal. The function $\mathbf{w}_\ell^2(\mathbf{x})$, coincides with the weight function, associated with the exceptional Laguerre polynomials [1]. Therefore, apart from an overall constant, $P_N(x)$ are the exceptional Laguerre polynomials $\hat{P}_{\ell,n}(x^2; g)$ with degree $2n + 2\ell$ in x [1]. This implies $N = 2(n + \ell)$, which when substituted in (19) gives the energy eigenvalue expression to be

$$E = 4n.\tag{23}$$

With $N = 2(n + \ell)$, it is easy to verify that (20), coincides with the differential equation [2] for the exceptional X_ℓ Laguerre polynomials

$$\begin{aligned}\partial_{xx}P_N(x) + 2 \left(-x + \frac{g + \ell}{x} - \frac{\partial_x \xi_\ell(x^2; g)}{\xi_\ell(x^2; g)} \right) \partial_x P_N(x) \\ \left(4n - 4\ell + 4x \frac{\partial_x \xi_\ell(x^2, g + 1)}{\xi_\ell(x^2, g)} \right) P_N(x) = 0.\end{aligned}\tag{24}$$

In addition, substituting the meromorphic form of the momentum function from (16) in (1), we obtain the unnormalized wave function as

$$\Psi_n(x) = \frac{x^{(g+\ell)} \exp(-\frac{1}{2}x^2)}{\xi_\ell(x^2; g)} \hat{P}_{\ell,n}(x^2; g),\tag{25}$$

which agrees with the known results [1]. Thus, we have obtained the energy eigenvalues and eigenfunctions for the new exactly solvable potentials, using the singularity structure of the momentum function. In addition to the fixed poles, q_ℓ has $2n$ moving poles on the real line and 2ℓ moving poles off the real line. The point at infinity turns out to be an isolated singularity.

Singularity structure

Here, we compare the singularity structure of the momentum function associated with these models to those of the QES and the conventional ES models. We see that the differences among these three models are due to the moving pole structures. For all the models, the number of moving poles located on the real line is consistent with the oscillation theorem. For the conventional ES models there were no moving poles off the real line. For the QES models, moving poles off the real line were found. The number of these poles varied from one QES level to the other, keeping the total number of moving poles (located at both real and complex locations) fixed for all solvable states. Compared to this, the new ES models have moving poles off the real line, but their number is fixed, equal to 2ℓ , for all the energy levels. Thus, we see that the singularity structure of the newly constructed potentials is neither completely like the ES models nor the QES models.

3 Exactness of SWKB quantization condition

In an earlier paper [15], the information on the singularity structure of the QMF for several models, was used to account for the exactness of the SWKB quantization condition,

$$\frac{1}{\pi} \int_{x_1}^{x_2} \sqrt{[E - W^2(x)]} dx = n, \quad (26)$$

where x_1 and x_2 ($x_1 < x_2$) are the turning points, which are the real roots of $\sqrt{E - W^2(x)}$.

For simplicity we denote the integrand $\sqrt{E - W^2(x)}$ as p_{SWKB} and put $\hbar = 1$. The integrals such as (26) can be turned into a contour integral (see for example [20]), and for (26) we get

$$I_C \equiv \frac{1}{2\pi} \oint_C p_{SWKB} dx = n, \quad (27)$$

where C is the contour enclosing the branch cut of p_{SWKB} from x_1 to x_2 . This integral can be computed in terms of the contributions of the singularities outside the contour C . The residues of p_{SWKB} are double valued and the right values are picked using the boundary condition, in the limit $E \rightarrow 0$, $p_{SWKB} \rightarrow p$, since $W(x) = -\psi'_0(x)/\psi_0(x)$. For the cases studied earlier in [15], the residues and the locations of the singularities outside the contours C of the two integrals, (27) and (2) matched identically and hence application of complex integration techniques gave same answers.

In the present case, we first list out the location of the poles of p_{SWKB} outside the contour C and corresponding residues in table 1 below. It can be easily seen that the

| location | residue |
|---|------------------------------|
| $x = 0$ | $-i(g + \ell)$ |
| 2ℓ singularities located at the zeros of $\xi_\ell(x^2, g + 1)$ | $-i$ |
| 2ℓ singularities located at the zeros of $\xi_\ell(x^2, g)$ | i |
| $x = \infty$ | $\frac{iE}{2} + i(g + \ell)$ |

Table 1: Pole structure of p_{SWKB} and $p(x)$.

poles and their residues are identical to those of the QMF. However in this case, p_{SWKB} has additional branch points as illustrated in figure 1, for the case of $l = 1$ and $g = 1$. In figure 1, the contour Θ encloses all the poles and branch cuts of p_{SWKB} . The contour C_1 encloses the origin and the contour C_2 encloses the branch cut between $-x_1$ and $-x_2$, which is equal to the integral in (27) by symmetry. The contours γ_m, Γ_m enclose the simple poles corresponding to the 2ℓ zeros of $\xi_\ell(x^2, g)$ and $\xi_\ell(x^2, g + 1)$ respectively. Finally, the contours Ω_i enclose the branch cuts of p_{SWKB} located off the real line. By writing the left hand side of (27) in terms of contour integrals enclosing the poles and




Figure 1: The figure displays the singularities and branch cuts of $\sqrt{E - W^2(x)}$, for $g = 1$ and $l = 1$ and the contours enclosing them. All the contours here are oriented in the anti-clockwise direction.

branch cuts of p_{SWKB} located outside C gives

$$\begin{aligned}
 I_C = & \frac{1}{2\pi} \oint_{\Theta} p_{SWKB} dx - \frac{1}{2\pi} \left(\oint_{C_1} p_{SWKB} dx + \oint_{C_2} p_{SWKB} dx \right. \\
 & \left. + \sum_{m=1}^{2\ell} \left[\oint_{\gamma_m} p_{SWKB} dx + \oint_{\Gamma_m} p_{SWKB} dx \right] + \sum_{i=1}^{4\ell} \oint_{\Omega_i} p_{SWKB} dx \right). \quad (28)
 \end{aligned}$$

We show that the contribution of the branch cuts of p_{SWKB} enclosed by contours Ω_i should vanish, for the SWKB condition to be exact. For this purpose we will evaluate the different integrals, except the ones around the contours Ω_i in the left hand side of (28). The integral around contour Θ is equal to $2\pi i$ (residue at x equal to ∞), calculated using the rule [20]

$$\begin{aligned} \text{Residue of } \sqrt{E - W^2(x)} \quad \text{at } z \text{ equal to } \infty &= - \text{Coefficient of } t \text{ in the} \\ &\text{expansion of } \sqrt{E - W^2(1/t)} \text{ for small } t. \end{aligned} \quad (29)$$

By using $\frac{\partial_x \xi_\ell(x^2; g+1)}{\xi_\ell(x^2; g+1)} = \sum_{i=1}^{2\ell} \frac{1}{x-a_i}$ and $\frac{\partial_x \xi_\ell(x^2; g)}{\xi_\ell(x^2; g)} = \sum_{j=1}^{2\ell} \frac{1}{x-b_j}$, one obtains

$$\sqrt{E - W^2(1/t)} = \frac{1}{t} \sqrt{Et^2 - \left(1 - (g + \ell)t^2 - t \sum_{i=1}^{2\ell} \frac{t}{1 - ta_i} + t \sum_{j=1}^{2\ell} \frac{t}{1 - tb_j} \right)^2}, \quad (30)$$

$$\equiv \pm \frac{i}{t} \sqrt{1 - Et^2 - 2(g + \ell)t^2 - \sum_{i=1}^{2\ell} \frac{t^2}{1 - ta_i} + \sum_{j=1}^{2\ell} \frac{t^2}{1 - tb_j} + H(t)}, \quad (31)$$

where $H(t)$ denotes all the other higher order terms. Expanding the square root, one obtains

$$\sqrt{E - W^2(1/t)} \equiv \pm \frac{i}{t} \left(1 - \frac{1}{2}(E + 2(g + \ell)^2)t^2 + \dots \right), \quad (32)$$

which gives the residue at infinity to be $\pm i(\frac{E}{2} + (g + \ell))$. The application of the boundary condition $E \rightarrow 0$, $p_{SWKB} \rightarrow p$, allows us to choose the right value of the residue to be $i(\frac{E}{2} + (g + \ell))$. Therefore

$$\begin{aligned} \frac{1}{2\pi} \oint_{\Theta} p_{SWKB} dx &= -i \left(i \frac{E}{2} + i(g + \ell) \right) \\ &= \left(\frac{E}{2} + (g + \ell) \right) \end{aligned} \quad (33)$$

The other method to calculate the integral around Θ is to use the mapping $x = 1/t$, which maps the point at infinity in the x plane to the origin in the t plane. The coefficient of $1/t$ in the Laurent expansion of the integrand in powers of t , will be the residue and is same as the value calculated above.

The residues at the other singular points, listed in table 1, can be obtained by using simple complex variable techniques. Now with all the residues known, application of the Cauchy residue theorem gives,

$$I_C = \frac{E}{2} + (g + \ell) - \left((g + \ell) + I_C + [-2\ell + 2\ell] + \frac{1}{2\pi} \sum_{i=1}^{4\ell} \oint_{\Omega_i} p_{SWKB} dx \right). \quad (34)$$

Simplifying the right hand side gives

$$2I_C = \frac{E}{2} - \frac{1}{2\pi} \sum_{i=1}^{4\ell} \oint_{\Omega_i} p_{SWKB} dx. \quad (35)$$

If E coincides with the n^{th} eigenvalues given in (23), we obtain

$$I_C = n - \frac{1}{4\pi} \sum_{i=1}^{4\ell} \oint_{\Omega_i} p_{SWKB} dx. \quad (36)$$

Therefore, it is obvious that if the sum of the contribution of the branch cuts off the real line vanishes, we obtain

$$\frac{1}{2\pi} \oint_C p_{SWKB} = n \quad (37)$$

Thus vanishing of the contribution of the branch cuts off the real line turns out to be a necessary and sufficient condition for the exactness of the SWKB rule.

We point out here that the calculation of the integral appearing in (2), in terms of p , also proceeds in a similar fashion. The only difference being that the QMF does not have branch points or moving branch point singularities in the complex plane. Therefore this integral receives contribution only from the fixed poles corresponding to the potential, the n moving poles located symmetrically between $-x_1$ and $-x_2$ and from the point at infinity. As the table suggests the contribution from these poles, for both integrals is identical. Therefore it is obvious that the SWKB condition is exact if the contribution from the branch cuts off the real line vanishes, when the energy is an eigenvalue.

In [16],[21] it has been proved that the SWKB condition is exact for all the conventional ES, translationally SIPs. The proof makes use of the property of shape invariance, where the assumption that $W^2(x)$ is of $O(\hbar^0)$ while $\hbar\partial_x W(x)$ is of $O(\hbar)$ was the key point. However, unlike the conventional SIPs with translation, for the new SIPs with translation, $W(x)$ is in fact a complicated function of \hbar . Thus the derivation in [16], [21] about the exactness of SWKB for SIPs with translation is strictly not applicable for the newly discovered SIPs with translation. Our present study of the exactness of the SWKB rule suggests that the disappearance of the contribution of the branch cuts off the real line, when E is equal to the one of the eigenvalue is possibly linked to the property of shape invariance.

4 Conclusions

In this study, we have analyzed the new ES shape invariant potentials obtained by deforming the radial oscillator using the QHJ formalism and obtained the expressions for the eigenvalues and eigenfunctions. We have investigated the singularity structure of the QMF and compared it with those of the conventional ES and QES models studied earlier. The QMF for these potentials has $2n$ real and 2ℓ complex moving poles, with $E = 4n$ giving the energy eigenvalue and ℓ fixing the potential. We found that the singularity structure of these models is neither completely like the ES models nor the QES models. The exactness of the SWKB rule has been shown to be equivalent to vanishing of the contribution of the branch cuts off the real line. Lastly, we point out that the above results can be extended to the other families of infinite number of potentials obtained by deforming the Darboux-Pöschl-Teller potentials [1].

Acknowledgments S S R thanks the Centre for Advanced Studies (CAS), School of Physics, University of Hyderabad, Hyderabad and the Department of Science and Technology (DST), India (fast track scheme (D. O. No: SR/FTP/PS-13/2009)) for financial support.

5 References

References

- [1] Odake S and Sasaki R 2009 *Phys. Lett. B* **679** 414 (arXiv:0906.0142).
- [2] Sasaki R, Tsujimoto S and Zhedavov A 2010 *J. Phys. A: Math. Theor* **43** 315204 (arXiv:1004.4711).
- [3] Darboux G, Acad C R 1882 *Paris* **94** 1456.
- [4] Crum M M 1955 *Quart. J. Math. Oxford Ser.(2)* **6** 121 (arXiv:9908019).
- [5] Gómez-Ullate D, Kamran N and Milson R 2010 *J Approx Theory* **162** 987 (arXiv:0805.3376).
- [6] Gómez-Ullate D, Kamran N and Milson R 2009 *J. Math. Anal. Appl.* **359** 352 (arXiv:0807.3939).
- [7] Leacock R A and Padgett M J 1983 *Phys. Rev. D* **28** 2491.
- [8] Leacock R A and Padgett M J 1983 *Phys. Rev. Lett.* **50** 3.
- [9] Bhalla R S, Kapoor A K and Panigrahi P K 1997 *Am. J. Phys.* **65** 1187.
- [10] Sree Ranjani S, Geojo K G, Kapoor A K and Panigrahi P K 2004 *Mod. Phys. Lett. A* **19** 1457.
- [11] Geojo K G , Sree Ranjani S and Kapoor A K 2003 *J. Phys. A: Math. Gen.* **36** 4591.
- [12] Sree Ranjani S 2005 *Quantum Hamilton - Jacobi solution for spectra of several one dimensional potentials with special properties* Thesis submitted to the University of Hyderabad (arXiv:0408036).
- [13] Dutt R, Khare A and Sukhatme U P 1988 *Am. J. Phys.* **56** 163.
- [14] Raghunathan K, Seetharaman M and Vasani S S 1987 *Phys. Lett. B* **188** 351.
- [15] Bhalla R S, Kapoor A K and Panigrahi P K 1996 *Phys. Rev. A* **54** 951.
- [16] Cooper F, Khare A and Sukhatme U P 2001 *Supersymmetric quantum mechanics*(Singapore: World Scientific Publishing Co. Ltd.)
- [17] Quesne C 2008 *J. Phys. A: Math. Theor.* **41** 392001.
- [18] Bougie J, Gangopadhyaya A, and Mallow J V 2010 *Phys. Rev. Lett.* **105** 210402.
- [19] Morse P M, Feshbach H 1953 *Methods of theoretical physics (part 1)* (New York: McGraw-Hill Book Company Inc.) pp. 784-785.
- [20] Kapoor A K 2011 *Complex variables, principles and problem sessions* (Singapore: World Scientific Publishing Co. Ltd.) chap. 6, sec. 8.
- [21] Dutt R, Khare A, Sukhatme U P 1986 *Phys. Lett. B* **181** 295.