



10-25-2010

# Synthesis of an Ortho-Triazacyclophane: N,N',N''-Trimethyltribenzo-1,4,7-triazacyclononatriene

Andria M. Panagopoulos  
*Oak Ridge National Laboratory*

Matthias Zeller  
*Youngstown State University*

Daniel P. Becker  
*Loyola University Chicago, dbecke3@luc.edu*

## Author Manuscript

This is a pre-publication author manuscript of the final, published article.

## Recommended Citation

Panagopoulos, Andria M.; Zeller, Matthias; and Becker, Daniel P. Synthesis of an Ortho-Triazacyclophane: N,N',N''-Trimethyltribenzo-1,4,7-triazacyclononatriene. *Journal of Organic Chemistry*, 75, 22: , 2010. Retrieved from Loyola eCommons, Chemistry: Faculty Publications and Other Works, <http://dx.doi.org/10.1021/jo1017074>

This Article is brought to you for free and open access by the Faculty Publications at Loyola eCommons. It has been accepted for inclusion in Chemistry: Faculty Publications and Other Works by an authorized administrator of Loyola eCommons. For more information, please contact [ecommons@luc.edu](mailto:ecommons@luc.edu).



This work is licensed under a [Creative Commons Attribution-Noncommercial-No Derivative Works 3.0 License](https://creativecommons.org/licenses/by-nc-nd/3.0/).

© 2010 American Chemical Society

**Synthesis of an ortho-Triazacyclophane: N,N',N''-Trimethyl  
Tribenzo-1,4-7-triazacyclononatriene**

Journal:	<i>The Journal of Organic Chemistry</i>
Manuscript ID:	jo-2010-017074.R1
Manuscript Type:	Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Panagopoulos, Andria; Loyola University, Chemistry Zeller, Matthias; Youngstown State University, Department of Chemistry Becker, Daniel; Loyola University, Department of Chemistry

SCHOLARONE™  
Manuscripts

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Synthesis of an *ortho*-Triazacyclophane: *N,N',N''*-Trimethyl tribenzo-1,4-7-triazacyclononatriene


Andria M. Panagopoulos<sup>§</sup>, Matthias Zeller<sup>†</sup> and Daniel P. Becker<sup>§\*</sup>

<sup>§</sup>Department of Chemistry, Loyola University Chicago, 1032 W. Sheridan Road,

Chicago, IL 60660. <sup>†</sup>Department of Chemistry, 1 University Plaza, Youngstown State

University, Youngstown, OH 44555-3663.

\*dbecke3@luc.edu



### Abstract


*N,N',N''*-Trimethyl tribenzo-1,4-7-triazacyclononatriene has been synthesized via sequential palladium-catalyzed Buchwald-Hartwig N-arylation reactions affording the 9-membered triaza *ortho*-cyclophane in 35% overall yield. An X-ray crystal structure shows the new cyclophane to have a C<sub>2</sub>-symmetric saddle conformation, as compared to the crown conformation exhibited by the related carbocyclic cyclotrimeratrylene (CTV).

## Introduction

The area of supramolecular chemistry is of continued interest due to a wide variety of applications including materials technology, catalysis, medicine, analytical detection and sensing. Cyclophanes, which are supramolecular structures comprised of aromatic units with bridging chains, have applications in molecular recognition as synthetic receptors<sup>1</sup> and have been used as building blocks for organic catalysts.<sup>2</sup> There is growing interest in the development of cyclophanes as hosts for ionic guests.<sup>3-5</sup>

The archetypal cyclophane cyclotrimeratrylene (CTV, **1**), a [1.1.1]orthocyclophane, is a commonly employed scaffold in supramolecular chemistry<sup>6</sup> that is readily prepared from veratryl alcohol in acid, and has been studied extensively for its capability of binding a number of smaller organic and organometallic guests within its bowl-shaped cleft.<sup>7-9</sup> CTV modification continues to be a significant area of study<sup>10-13</sup> and it has been used as a building block enabling the construction of more complex cryptophanes.<sup>14-17</sup> Although CTV has proven to be quite useful the molecule suffers from insolubility in aqueous systems and only rare opportunities for derivatization of the apical methylenes; most derivatives of CTV have been prepared by varying the groups on the phenolic oxygens around the periphery of the molecule; Collet was among the first to transform CTV into cryptophanes in this manner.<sup>18</sup> Nierengarten addressed the aqueous insolubility of CTV by appending polyethylene glycol units via the peripheral oxygens toward derivatives with biomedical applications, specifically to aid in the biological delivery of fullerenes, although the derivatives were of high molecular weight (>3000 to >6000 amu).<sup>19</sup> As part of our own exploration of apex-modified CTV derivatives<sup>20</sup> we have isolated the crown

1  
2  
3 and saddle conformers of CTV-oxime (**2a,b**)<sup>21</sup> and studied the kinetics and  
4  
5 thermodynamics of their interconversion between the crown and saddle CTV-oxime  
6  
7 derivative **2a,b**<sup>22</sup> (Figure 1). CTV exists almost exclusively in its crown conformation  
8  
9 (**1a**), and the saddle conformer of CTV was only recently isolated and characterized  
10  
11 through high temperature melt and quench techniques by Zimmermann who also studied  
12  
13 the thermodynamic and kinetic properties of the interconversion of the crown and saddle  
14  
15 CTV conformers.<sup>23</sup> Holman<sup>24</sup> as well as Huber<sup>25</sup> have identified topoisomeric  
16  
17 cryptophanes containing blended crown and saddle CTV moieties; Holman's  
18  
19 cryptophane undergoes a conformational crown to saddle "implosion" upon thermal  
20  
21 liberation of its tetrahydrofuran guest.  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33




**Figure 1.** Cyclotrimeratrylene (CTV) and CTV-oxime derivatives.

Several heteroatom analogues of CTV and the tribenzocyclononene core have been reported in which the methylene groups have been replaced (**3a-c**).

Trithiacyclotrimeratrylene (**3a**)<sup>26, 27</sup> forms complexes with copper(I),<sup>28</sup> rhodium (III),<sup>29</sup> and

1  
2  
3 platinum (II),<sup>30</sup> and it also exists in a temperature and solvent-dependent equilibrium of  
4  
5 the crown and saddle forms. In addition, the trioxycyclononene **3b**,<sup>31, 32</sup> and trimercury  
6  
7 **3c**,<sup>33</sup> which is planar rather than crown-shaped have also been described. Recently, the  
8  
9 first apical methylene aza-substituted cyclophane was reported with the synthesis of  
10  
11 monoamine tribenzo-1-azacyclophane **3d**<sup>34</sup> which was prepared as a potential  
12  
13 benzodiazepine receptor ligand, while Tanaka reported the synthesis of the trimethyl  
14  
15 triaza[1<sub>3</sub>]metacyclophane **4**.<sup>35</sup> We envisioned a derivative of the  
16  
17 tribenzotricyclononatriene core of CTV retaining its high (C<sub>3v</sub>) symmetry with nitrogen  
18  
19 atoms in place of the methylenes, to provide a ready handle for apical functionalization  
20  
21 and altering the charge on the scaffold, to enable modulation of host-guest properties and  
22  
23 to facilitate attachment to surfaces. Furthermore, the nitrogen atoms should function as a  
24  
25 ligand for metals as described for the oxa- and thia-analogs above, ultimately providing a  
26  
27 redox-switchable host molecule. Herein we describe the synthesis of the novel triaza-  
28  
29 *orthocyclophane* **5a** (Figure 2).  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




**Figure 2.** Analogs of CTV

## Results and Discussion

Looking retrosynthetically at the target tribenzo-1,4,7-triazacyclononatriene, we envisioned four unique synthetic approaches (Figure 3), with N-aryl bonds constructed potentially utilizing Buchwald-Hartwig<sup>36-39</sup>, Ullman<sup>40</sup>, or nucleophilic aromatic substitution ( $S_{\text{N}}\text{AR}$ ) methodologies. Route A considers the direct trimerization of an ortho-substituted aniline. This highly convergent synthesis has literature precedent in the work of Tanka<sup>35</sup> and coworkers who synthesized *meta*-cyclophane **4** in low yield directly from a *meta*-substituted aniline. Approaches B and C consider a diphenylamine derivative joining the third ring in a double-coupling. Finally, route D describes a linear construction of the molecule. The final intramolecular macrocyclization of the 9-membered ring should be aided by the limited degrees of freedom imposed by the aryl

rings (Figure 3).




**Figure 3.** Retrosynthetic strategies toward the tribenzo-1,4-7-triazacyclononatriene ring system

For the highly-convergent approach (A), we subjected 2-iodoaniline and 2-bromoaniline to a variety of Buchwald-Hartwig cross-coupling conditions. Employing Pd(OAc)<sub>2</sub>, P(*t*-Bu)<sub>3</sub> as the ligand, and sodium tert-butoxide as the base in dioxane under reflux as a typical example afforded only phenazine in 92% isolated yield, via facile air oxidation of the metastable dihydrophenazine.<sup>41</sup> Tanaka was able to synthesize the triaza-*metacyclophane* **4** from the corresponding secondary *N*-methyl derivative but not from the unsubstituted (primary) aniline. Therefore we attempted the convergent trimerization employing a secondary aniline utilizing similar Pd-catalyzed cross coupling conditions. *N*-Methyl-2-bromoaniline which under Buchwald-Hartwig conditions produced a



1  
2  
3 complex mixture from which only dehalogenated dimer and dehalogenated tetramer were  
4  
5 isolated in extremely low yields (0.3% and 2%, respectively). We were unable to detect  
6  
7 the desired product in any mixture derived from the direct trimerization approach under a  
8  
9 variety of Buchwald-Hartwig conditions, even when an authentic sample was available  
10  
11 for direct comparison from the linear approach below.  
12  
13  
14  
15  
16  
17


18 Given the facile formation of phenazine in the trimerization attempt, we focused on the  
19  
20 moderately convergent [2x1] syntheses (Routes B and C). Subjecting bis-(2-  
21  
22 bromophenyl)amine to Buchwald-Hartwig conditions, carbazole **6** was formed as the  
23  
24 major product via a reductive coupling in 65% yield. In addition, the phenazine  
25  
26 derivative **7** was produced in 4% via intramolecular *N*-arylation involving a 6-member  
27  
28 ring formation. The mono-coupled triaryl product **8** was also isolated in 4% yield  
29  
30  
31 (Scheme 1).  
32  
33  
34  
35



50 **Scheme 1.** 1 x 2 synthesis attempt from *bis*-(2-bromophenyl)amine  
51  
52  
53

54 A similar convergent synthesis beginning with *N*-(2-aminophenyl)-1,2-benzenediamine<sup>42</sup>,  
55  
56  
57 <sup>43</sup> is shown in scheme 4. The diamine and the 1,2-dihalide were subjected to palladium-  
58  
59  
60


1  
2  
3 catalyzed N-arylation producing phenazine **9** in 20% yield as the only isolated product.  
4  
5  
6 When an N-methyl blocking group was added<sup>44</sup> to block the central nitrogen, attempted  
7  
8 N-arylation of the *N*-(2-aminophenyl)-*N*-methyl-1,2-benzenediamine with 1,2-  
9  
10 dibromobenzene resulted in *N,N'*-dimethyl phenazine derivative **10** as the only isolated  
11  
12 product (Scheme 2).  
13  
14  
15  
16  
17  
18  
19



32 **Scheme 2.** 1 x 2 synthesis attempt from *N*-(2-aminophenyl)-1,2-benzenediamine  
33  
34  
35  
36

37 The inability to produce the desired cyclophane utilizing convergent approaches led us to  
38 pursue a linear synthesis (Route D) with sequential protection of the aniline nitrogens to  
39 avoid 6-membered ring formation. Scheme 5 outlines the successful synthesis of the  
40  
41 *N,N',N''*-trimethyl triaza-*orthocyclophane* **5a** in 35% overall yield. A modification of  
42  
43 the Buchwald-Hartwig N-arylation method used by Tietze<sup>42</sup> employing 1,2-  
44  
45 bromochlorobenzene and *o*-nitroaniline gave 2-chloro-*N*-(2-nitrophenyl)-benzenamine **11**  
46  
47 in 99% yield. The aniline was protected by methylation using KOH and Me<sub>2</sub>SO<sub>4</sub> in  
48  
49 refluxing acetone<sup>45</sup> to give the *N*-methyl diphenylamine **12** in quantitative yield without  
50  
51 the need for further purification. Compound **12** was easily reduced with the general  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 method of Sanz<sup>46</sup> using CuCl and KBH<sub>4</sub> in dry MeOH at room temperature to give **13** in  
4  
5 quantitative yield, again without further purification, whereas more common reduction  
6  
7 methods such as Pd/C and H<sub>2</sub>NNH<sub>2</sub> or hydrogenation gave dehalogenated or  
8  
9 demethylated reduction products. The aniline **13** was coupled to iodonitrobenzene using  
10  
11 the previously established Buchwald-Hartwig conditions to produce the triaryl amine **14**  
12  
13 in 80% yield. Compound **14** was subjected to methylation by KH and MeI followed by  
14  
15 reduction of the nitro group with CuCl and KBH<sub>4</sub> to give products **15** and **16**,  
16  
17 respectively. The triaryl aniline was successfully closed to the 9-membered cyclophane  
18  
19 through the use of Buchwald-Hartwig coupling in a microwave reactor to afford the  
20  
21 *N,N'*-dimethyl triaza-*orthocyclophane* **5b** in 50% isolated yield after purification. The  
22  
23 last step is a macrocyclization with unique steric demands and application of thermal  
24  
25 conditions gave significant decomposition, low yields (<5%), poor conversion and  
26  
27 extremely long reaction times (> 5 days). However we found that application of  
28  
29 microwave conditions in this step gave moderate to good yields. The third apical  
30  
31 nitrogen was methylated employing KH and MeI to give a quantitative yield of the  
32  
33 *N,N',N''*-trimethyl triaza-*orthocyclophane* **5a** without the need for further purification  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



**Scheme 3.** Reaction conditions: a. Pd(dba)<sub>2</sub>, BINAP, toluene, 24 h; b. KOH, Me<sub>2</sub>SO<sub>4</sub>, acetone; c. CuCl, KBH<sub>4</sub>, MeOH; d. KH, MeI, DMF; e. Pd(dba)<sub>2</sub>, BINAP, 1:5 *t*-BuOH:tol

The structure of the *N,N'*-dimethyl derivative **5b** was assigned based on the equivalent methyls in the <sup>1</sup>H NMR at δ 2.68 (6H, s) and the exact mass (MH<sup>+</sup>) observed by mass spectrometry. The structure was ultimately confirmed by single crystal X-ray analysis revealing a C<sub>2</sub>-symmetric saddle (Figure 4). The <sup>1</sup>H NMR of the *N,N',N''*-trimethyl derivative **5a** reveals the high C<sub>3v</sub> symmetry with chemical shifts δ 6.90 (12H, s), δ 2.91 (9H, s) manifesting the similarity of protons ortho and meta to the nitrogen, leading to a fortuitous singlet for all 12 aromatic protons.




Figure 4. X-ray structure of compound **5b** with atom labels, thermal ellipsoids are at the 50% probability level.

In conclusion, we have constructed the new triazacyclopentane, tribenzo-1,4,7-triazacyclononatriene **5b**, in 7 steps via palladium catalyzed C-N amination, followed by alkylation and reduction, and reiteration of this sequence in order to obtain the triaryl precursor to the final palladium-catalyzed cyclization to the 9-membered cyclophane. Alkylation of **5b** gives the  $C_{3v}$ -symmetric  $N,N',N''$ -trimethyl triaza-*orthocyclophane* and demonstrates the ability to functionalize the cyclophane at the apex in order to modulate its physicochemical properties. We envision that the new triazacyclopentane should will complement the familiar carbocyclic framework of CTV with greater versatility, including the ability to bind cationic species in the apex analogous to metal complexes of the well known triazacyclononene (TACN) derivatives that have utility as MRI contrast agents<sup>47, 48</sup> and radioimmunotherapy agents.<sup>49</sup> These studies are in progress and will be reported in due course.

Experimental

## General Experimental

All solvents were distilled prior to use. All reagents were used without further purification unless otherwise noted. All Pd-catalyzed and Cu-catalyzed reactions were conducted under an inert atmosphere of argon, and all other reactions were conducted under a nitrogen atmosphere. Sorbent Technologies silica gel 60A, 40–75  $\mu\text{m}$  (200 x 400 mesh) was used for column chromatography. Sorbent Technologies aluminum-backed Silica gel 200 $\mu\text{m}$  plates were used for TLC.  $^1\text{H}$  NMR spectra were obtained utilizing a 300 MHz spectrometer with trimethylsilane (TMS) as the internal standard.  $^{13}\text{C}$  NMR spectra were obtained using a 75 MHz spectrometer. A CEM Discover<sup>®</sup> Microwave Model # 908005 was used for all microwave (MW) reactions. Infrared (IR) spectra were determined as a solution in  $\text{CHCl}_3$ . Single crystal X-ray diffraction data were collected on a charge-coupled-device (CCD) diffractometer with a liquid nitrogen vapor cooling device. Data were collected at 100 K with a graphite monochromatized  $\text{MoK}\alpha$  X-ray radiation ( $\lambda = 0.71073 \text{ \AA}$ ). Data were collected and reduced and corrected for absorption using multi-scan methods. The structure was solved by direct methods and refined by full matrix least squares against  $F^2$  with all reflections. Non hydrogen atoms were refined anisotropically. C-H hydrogen atom positions were idealized. Additional details of the structure determination can be found in the supplementary cif file.

**(2'-Chlorophenyl)-(2-nitrophenyl)-amine (11)**. Compound **11** was synthesized according to the general procedures outlined by Tietze *et al.*<sup>42</sup> A pressure tube was charged with o-nitroaniline ( 0.690 g, 5 mmol), o-bromochlorobenzene ( 0.60 mL, 5.00

mmol), Pd(dba)<sub>2</sub> (0.144 g, 5%), BINAP (0.233 g, 7.5%), Cs<sub>2</sub>CO<sub>3</sub> (3.26 g, 10 mmol) and toluene (10 mL). The mixture was purged with argon for 10 min at rt and the pressure tube was sealed. The reaction was sealed and placed in a pre-heated oil bath. The temperature was brought to 120°C and the reaction stirred for 24 h. TLC showed complete consumption of o-nitroaniline and the reaction mixture was filtered through a pad of SiO<sub>2</sub> using 5/5/90 EA/DCM/petroleum ether as the eluent. The solvent was removed under vacuum and no further purification was needed to give the product as an orange solid (1.24 g, 100%) which was identical to the material reported in the literature<sup>42</sup> by <sup>1</sup>H NMR.

**2-Chloro-N-methyl-N-(2-nitrophenyl)aniline (12).** Following the general method of Wilshire<sup>45</sup> compound **11** ( 1.25 g, 5 mmol) was stirred at rt in acetone (16 mL) and freshly crushed KOH ( 1.23 g, 22.0 mmol) was added to the stirring mixture. After the reaction was brought to reflux, Me<sub>2</sub>SO<sub>4</sub> (2.18 mL, 23 mmol) was added dropwise via syringe over 10 min. The mixture was allowed to stir at reflux for 1 h. The reaction was cooled to rt and 20 mL of 10 M NaOH was added to the solution. After 1 h the mixture was quenched with 10 mL H<sub>2</sub>O and extracted 3 x 10 mL DCM. The organic layers were combined and dried over Na<sub>2</sub>SO<sub>4</sub>. The solvent was removed under vacuum and the mixture was placed in an 80°C oil bath under vacuum to remove excess Me<sub>2</sub>SO<sub>4</sub>. No further purification was needed to obtain the desired as an off-white solid (1.31 g, 100%). mp 73-75 °C; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.68 (1H, dd, J= 8.1, 1.5 Hz), 7.54 (1H, ddd, J= 8.7, 7.3, 1.7 Hz), 7.42 (1H, dd, J= 7.8, 1.5 Hz), 7.19 (1H, dd, J=7.7, 1.7 Hz), 7.14-7.11 (2H, m), 7.06 (1H, dd, J=7.7, 1.9 Hz), 7.0 (1H, ddd, J=8.2, 7.3, 1.1 Hz); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>) δ 145.0, 143.0, 133.2, 131.3, 130.9, 128.7, 127.9, 126.7, 126.2, 125.9,

1  
2  
3 120.8, 120.6, 41.1; IR (CDCl<sub>3</sub>): 1520 (NO<sub>2</sub>); HRMS (MH<sup>+</sup>) calcd for C<sub>13</sub>H<sub>11</sub>O<sub>2</sub>N<sub>2</sub>Cl  
4  
5 263.0509, found 263.0604.  
6  
7

8  
9  
10 **N<sup>1</sup>-(2-Chlorophenyl)-N<sup>1</sup>-methylbenzene-1,2-diamine (13)**. Following the general  
11 procedure of Sanz<sup>46</sup>, CuCl (0.137 g, 1.38 mmol) was added to a stirring solution of  
12 compound **12** (0.121 g, 0.46 mmol) in MeOH (4.6 mL) at rt. KBH<sub>4</sub> (0.174 g, 3.22 mmol)  
13 was then added in portions. The reaction stirred at rt until the solution became clear, (2-4  
14 h). The reaction was quenched with H<sub>2</sub>O and extracted 3 x 15 mL 90/10 EA/DCM. The  
15 organic layers were combined and dried over Na<sub>2</sub>SO<sub>4</sub> and the solvent was removed to  
16 give the desired product as a brown oil (0.107 g, 100%). <sup>1</sup>H NMR (300MHz, CDCl<sub>3</sub>) δ  
17 7.32 (1H, dd, J= 7.8, 1.4 Hz), 7.25 (1H, dd, J= 7.3, 1.7 Hz), 7.22 (1H, dd, J= 7.1, 1.7 Hz),  
18 7.16 (1H, dd, J= 8.0, 1.6 Hz), 7.00-6.95 (2H, m), 6.76 (1H, ddd, J=9.3, 7.7, 1.4 Hz), 6.67  
19 (1H, ddd, J= 8.9, 7.6, 1.5 Hz); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>) δ 147.6, 142.2, 136.9, 130.7,  
20 130.68, 127.4, 125.5, 123.6, 121.9, 118.6, 115.8, 41.1; IR (CDCl<sub>3</sub>): 3440 (NH<sub>2</sub>), 3351  
21 (NH<sub>2</sub>); HRMS (MH<sup>+</sup>) calcd for C<sub>13</sub>H<sub>13</sub>N<sub>2</sub>Cl 233.0767, found 233.0791.  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40

41 **N<sup>1</sup>-(2-Chlorophenyl)-N<sup>1</sup>-methyl-N<sup>2</sup>-(2-nitrophenyl)benzene-1,2-diamine (14)**.

42 Compound **13** (0.842 g, 3.62 mmol), o-iodonitrobenzene (1.35 g, 5.43 mmol), Pd(dba)<sub>2</sub>  
43 (0.104 g, 5% mol), BINAP (0.170 g, 7.5%), Cs<sub>2</sub>CO<sub>3</sub> (2.35 g, 7.42 mmol) and 12 mL of  
44 toluene were placed in a pressure tube. The mixture was purged with argon at rt for 15  
45 min and the tube was then sealed and placed in a pre-heated oil bath at 80-90°C for 30 h.  
46 After TLC showed consumption of **13**, the reaction mixture was filtered through a pad of  
47 silica gel eluting with 90/10 EA/DCM. The solvent was then removed under vacuum.  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 The crude product was then purified by column chromatography on silica gel eluting with  
4  
5 1/99 Et<sub>2</sub>O/petroleum ether to afford the desired product as a red crystalline solid (0.785 g,  
6  
7 80%): mp 141-145 °C; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 9.03 (1H, bs), 8.07 (1H, dd, J=8.7,  
8  
9 1.5 Hz), 7.32-7.19 (4H, m), 7.12-6.99 (5H, m), 6.90 (1H, ddd, J= 8.0, 6.9, 2.2 Hz), 6.68  
10  
11 (1H, ddd, J= 8.4, 6.9, 1.2 Hz), 3.16 (3H, s); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>) δ 147.2, 145.1,  
12  
13 142.4, 135.2, 131.6, 130.7, 129.5, 127.4, 126.5, 126.5, 126.0, 124.8, 124.0, 123.2, 121.7,  
14  
15 117.0, 115.8, 40.6; IR (CDCl<sub>3</sub>) 3344 (NH), 1503 (NO<sub>2</sub>); HRMS (MH<sup>+</sup>) calcd for  
16  
17 C<sub>19</sub>H<sub>16</sub>N<sub>3</sub>O<sub>2</sub>Cl 354.1009, found 354.0961.  
18  
19  
20  
21  
22  
23

24  
25 **N<sup>1</sup>-(2-Chlorophenyl)-N<sup>1</sup>,N<sup>2</sup>-dimethyl-N<sup>2</sup>-(2-nitrophenyl)benzene-1,2-diamine (15).** A  
26  
27 solution of compound **14** (0.405 g, 1.14 mmol) in 4 mL of DMF was added to KH (0.46  
28  
29 g, 3.42 mmol). Upon addition, the solution turned from orange to deep purple. The  
30  
31 mixture was stirred at rt for 10 min then MeI (0.4 mL, 5.7 mmol) was added dropwise via  
32  
33 syringe. The reaction was stirred at rt until the solution returned to a yellow color. The  
34  
35 reaction was then quenched with H<sub>2</sub>O and extracted 3 x 15 mL EA. The organic layers  
36  
37 were combined and washed 3 x 15 mL H<sub>2</sub>O, brine then H<sub>2</sub>O again to remove excess  
38  
39 DMF. The organic layer was then dried over MgSO<sub>4</sub>, the solvent was removed under  
40  
41 reduced pressure to give the desired product as a yellow powder with no further  
42  
43 purification necessary (0.362 g, 86%): <sup>1</sup>H NMR (300MHz, CDCl<sub>3</sub>) δ 7.63 (1H, dd, J=  
44  
45 8.0, 1.7 Hz), 7.36 (1H, ddd, J=8.8, 7.3, 1.8 Hz), 7.29-7.19 (2H, m), 7.12 (1H, dd, J=8.2,  
46  
47 1.7 Hz), 7.07-6.92 (5H, m), 6.88 (1H, ddd, J=8.2, 7.3, 1.2 Hz), 6.81 (1H, dd, J=7.8, 1.2  
48  
49 Hz), 3.32 (3H, s), 3.27 (3H, s); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>) δ 146.5, 143.5, 142.1, 138.8,  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 132.7, 131.0, 128.7, 127.6, 126.2, 124.2, 124.1, 123.7, 123.1, 120.4, 118.8, 38.5, 38.1; IR  
4  
5 (CDCl<sub>3</sub>) 1520 (NO<sub>2</sub>); HRMS (MH<sup>+</sup>) calcd for C<sub>20</sub>H<sub>18</sub>N<sub>3</sub>O<sub>2</sub>Cl 368.1166, found 368.1091.  
6  
7

8  
9  
10 **N<sup>1</sup>-(2-Aminophenyl)-N<sup>2</sup>-(2-chlorophenyl)-N<sup>1</sup>,N<sup>2</sup>-dimethylbenzene-1,2-diamine (16).**  
11

12 Following the general procedure of Sanz,<sup>46</sup> CuCl (0.460 g, 4.65 mmol) was added to a  
13 stirring solution of compound **15** (0.570 g, 1.55 mmol) in MeOH (15.5 mL) at rt. KBH<sub>4</sub>  
14 (0.836 g, 15.5 mmol) was then added in portions. The reaction stirred at rt until the  
15 solution became clear, 2-4 h. The reaction was then quenched with H<sub>2</sub>O and extracted 3  
16 x 30 mL 90/10 EA/DCM. The organic layers were combined and dried over Na<sub>2</sub>SO<sub>4</sub> and  
17 the solvent was removed to give the desired product as a reddish-brown oil, (0.450 g,  
18 86%): <sup>1</sup>H NMR (300MHz, CDCl<sub>3</sub>) δ 7.3 (1H, dd, J= 8.0, 1.9 Hz), 7.15 (1H, 8.5, 7.3, 1.7  
19 Hz), 7.06-6.88 (7H, m), 6.83-6.73 (2H, m), 6.60 (1H, dd, J=7.3, 1.5 Hz); <sup>13</sup>C NMR (75  
20 MHz, CDCl<sub>3</sub>) δ 147.0, 143.4, 142.2, 141.1, 136.7, 131.1, 128.2, 127.3, 124.6, 124.4,  
21 124.3, 123.4, 123.4, 123.2, 122.6, 122.5, 118.7, 116.1, 39.4, 38.6; IR (CDCl<sub>3</sub>) 3441  
22 (NH<sub>2</sub>), 3368 (NH<sub>2</sub>); HRMS (MH<sup>+</sup>) calcd for C<sub>20</sub>H<sub>20</sub>N<sub>3</sub>Cl 338.1424, found 338.1379.  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40

41 **N-Methyl-2-(10-methylphenazin-5(10H)-yl)aniline (5b).** Compound **16** (0.090 g, 0.27  
42 mmol), Pd(dba)<sub>2</sub> (0.016 g, 10% mol), BINAP (0.034 g, 20% mol), Cs<sub>2</sub>CO<sub>3</sub> (0.132 g, 0.41  
43 mmol) in 3 mL of 1:1 toluene/t-BuOH were added to a 10 mL microwave tube. The  
44 mixture was purged with Ar for 5 min while stirring at rt. The MW settings were as  
45 follows; P = 250W, Time = 60 min, Temp = 130°C; PSI = 250. The reaction mixture was  
46 checked by TLC after each 60 min run. When TLC showed consumption of starting  
47 material **16** (240 min total), the mixture was filtered through a pad of silica gel eluting  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 with 90/10 EA/DCM. The solvent was removed under reduced pressure. The product  
4  
5 was then purified by column chromatography eluting with DCM/petroleum ether gradient  
6  
7 to give the final product as a white powder (41mg, 50%):  $^1\text{H}$  NMR (300MHz,  $\text{CDCl}_3$ )  $\delta$   
8  
9 7.05-6.95 (6H, m), 6.87 (1H, d,  $J= 1.1$  Hz), 6.84 (1H, d,  $J= 1.1$  Hz), 6.75-6.73 (4H, m),  
10  
11 5.82 (1H, bs), 2.68 (6H, s);  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ )  $\delta$  144.2, 141.4, 140.2, 127.8,  
12  
13 126.1, 121.9, 121.1, 118.4, 117.5, 39.9; IR ( $\text{CDCl}_3$ ) 3382 (NH), 1499 (C=C); HRMS  
14  
15 (MH+) calcd for  $\text{C}_{20}\text{H}_{19}\text{N}_3$  302.1579, found 302.1573. Single crystals for X-ray  
16  
17 structural analysis were grown by the slow evaporation of DCM.  
18  
19  
20  
21  
22  
23  
24

25 **5,10,15-Trimethyl-10,15-dihydro-5H-tribenzo[b,e,h][1,4,7]triazonine (5a).** A  
26  
27 solution of dimethyl azacyclophane **5b** (0.018 g, 0.07 mmol) in 0.2 mL of DMF was  
28  
29 added to KH (0.028 g, 0.21 mmol). Upon addition, effervescence ensued and the  
30  
31 solution turned a pale pinkish-purple color. The mixture was stirred at rt until  
32  
33 effervescence ceased (about 5 min), and MeI (0.022 mL, 0.35 mmol) was added dropwise  
34  
35 via syringe. The reaction was allowed to stir at rt for 2 h, during which the solution  
36  
37 became faint yellow in color. The reaction mixture was quenched with deionized  $\text{H}_2\text{O}$   
38  
39 and extracted with EA (3 x 10 mL). The organic layers were combined and dried over  
40  
41  $\text{Na}_2\text{SO}_4$ , the solvent was removed under reduced pressure to give the product as a pale  
42  
43 yellow oil. (0.021 g, 95% yield):  $^1\text{H}$  NMR (300MHz,  $\text{CDCl}_3$ )  $\delta$  6.92 (12H, s), 2.93 (9H,  
44  
45 s);  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ )  $\delta$  144.0, 122.9, 121.0, 40.6; IR ( $\text{CDCl}_3$ ) 3052 (CH), 1605  
46  
47 (C=C) HRMS (MH+) calcd for  $\text{C}_{21}\text{H}_{21}\text{N}_3$  316.1808, found 316.1799.  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6 **Supporting Information Available:** X-ray crystal structure coordinates, and files for  
7  
8 compound **5b** in CIF format.  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra are available for all compounds in  
9  
10 the 8-step linear synthesis. This material is available free of charge via the Internet at  
11  
12 <http://pubs.acs.org>.  
13  
14  
15  
16  
17  
18  
19

## 20 **References**

- 21  
22  
23  
24 (1) Diederich, F. In *Cyclophanes*; The Royal Society of Chemistry: Cambridge, 1991; ,  
25  
26 pp 313.  
27  
28 (2) Vögtle, F. In *Cyclophane Chemistry: Synthesis, Structures and Reactions*; John  
29  
30 Wiley & Sons Ltd: West Sussex, England, 1993; , pp 501.  
31  
32 (3) Richard, J.; Pamart, M.; Hucher, N.; Jabin, I. *Tetrahedron Lett.* **2008**, *49*, 3848-3852.  
33  
34 (4) Conejo-Garcia, A.; Campos, J. M.; Sanchez-Martin, R. M.; Gallo, M. A.; Espinosa,  
35  
36 A. J. *Med. Chem.* **2003**, *46*, 3754-3757.  
37  
38 (5) Bruno, G.; Cafeo, G.; Kohnke, F. H.; Nicolo, F. *Tetrahedron* **2007**, *63*, 10003-10010.  
39  
40 (6) Collet, A. *Tetrahedron* **1987**, *43*, 5725-5759.  
41  
42 (7) Collet, A. *Comprehensive Supramolecular Chemistry* **1996**, *6*, 281-303.  
43  
44 (8) Steed, J. W.; Zhang, H.; Atwood, J. L. *Supramolecular Chemistry* **1996**, *7*, 37-45.  
45  
46 (9) Burlinson, N. E.; Ripmeester, J. A. *Journal of Inclusion Phenomena* **1984**, *1*, 403-  
47  
48 409.  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 (10) Nierengarten, J. *Fullerenes, Nanotubes & Carbon Nanostructures* **2005**, *13*, 229-  
4  
5 242.  
6  
7  
8 (11) Harig, M.; Neumann, B.; Stammeler, H.; Kuck, D. *European Journal of Organic*  
9  
10 *Chemistry* **2004**, 2381-2397.  
11  
12 (12) Hardie, M. J.; Ahmad, R.; Sumbly, C. J. *New Journal of Chemistry* **2005**, *29*, 1231-  
13  
14 1240.  
15  
16  
17 (13) Ahmad, R.; Hardie, M. J. *Supramolecular Chemistry* **2006**, *18*, 29-38.  
18  
19 (14) Collet, A.; Dutasta, J. P.; Lozach, B.; Canceill, J. *Top. Curr. Chem.* **1993**, *165*, 103-  
20  
21 129.  
22  
23  
24 (15) Matsubara, H.; Hasegawa, A.; Shiwaku, K.; Asano, K.; Uno, M.; Takahashi, S.;  
25  
26 Yamamoto, K. *Chem. Lett.* **1998**, 923-924.  
27  
28 (16) Matsubara, H.; Oguri, S.; Asano, K.; Yamamoto, K. *Chem. Lett.* **1999**, 431-432.  
29  
30 (17) Brotin, T.; Dutasta, J. *Chem. Rev. (Washington, DC, U. S. )* **2009**, *109*, 88-130.  
31  
32  
33 (18) Canceill, J.; Collet, A.; Gottarelli, G.; Palmieri, P. *J. Am. Chem. Soc.* **1987**, *109*,  
34  
35 6454-6464.  
36  
37  
38 (19) Rio, Y.; Nierengarten, J. *Tetrahedron Lett.* **2002**, *43*, 4321-4324.  
39  
40 (20) Lutz, M. R., Jr.; Zeller, M.; Becker, D. P. *Tetrahedron Lett.* **2008**, *49*, 5003-5005.  
41  
42  
43 (21) Lutz, M. R.; French, D. C.; Rehage, P.; Becker, D. P. *Tetrahedron Lett.* **2007**, *48*,  
44  
45 6368-6371.  
46  
47  
48 (22) David C. French; Marlon R. Lutz, J.; Chichi Lu; Matthias Zeller; and Daniel P.  
49  
50 Becker *J. Phys. Chem. A* **2009**, *113*, 8258.  
51  
52 (23) Zimmermann, H.; Tolstoy, P.; Limbach, H.; Poupko, R.; Luz, Z. The Saddle Form  
53  
54 of Cyclotrimeratrylene. *J Phys Chem B* **2004**, *108*, 18772-18778.  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4 (24) Mough, S. T.; Goeltz, J. C.; Holman, K. T. Isolation and Structure of an "Imploded"  
5  
6 Cryptophane. *Angew.Chem., Int.Ed.; Angewandte Chemie, International Edition*  
7  
8 **2004**, *43*, 5631-5635.  
9
- 10 (25) Huber, G.; Brotin, T.; Dubois, L.; Desvaux, H.; Dutasta, J.; Berthault, P. Water  
11  
12 Soluble Cryptophanes Showing Unprecedented Affinity for Xenon: Candidates as  
13  
14 NMR-Based Biosensors. *J. Am. Chem. Soc.* **2006**, *128*, 6239-6246.  
15  
16
- 17 (26) Weiss, T.; Klar, G. *Zeitschrift für Naturforschung, Teil B: Anorganische Chemie,*  
18  
19 *Organische Chemie* **1979**, *34B*, 448-450.  
20  
21
- 22 (27) Von Deuten, K.; Kopf, J.; Klar, G. *Crystal Structure Communications* **1979**, *8*, 569-  
23  
24 575.  
25  
26
- 27 (28) Von Deuten, K.; Kopf, J.; Klar, G. *Crystal Structure Communications* **1979**, *8*, 721-  
28  
29 728.  
30  
31
- 32 (29) Kopf, J.; Von Deuten, K.; Klar, G. *Crystal Structure Communications* **1979**, *8*,  
33  
34 1011-1016.  
35  
36
- 37 (30) Von Deuten, K.; Klar, G. *Crystal Structure Communications* **1981**, *10*, 757-764.  
38
- 39 (31) Weiss, T.; Schulz, P.; Klar, G. *Zeitschrift für Naturforschung, Teil B: Anorganische*  
40  
41 *Chemie, Organische Chemie, Biochemie, Biophysik, Biologie* **1974**, *29*, 156-158.  
42  
43
- 44 (32) Von Deuten, K.; Klar, G. *Zeitschrift für Naturforschung, Teil B: Anorganische*  
45  
46 *Chemie, Organische Chemie* **1981**, *36B*, 1526-1531.  
47
- 48 (33) Woodard, C. M.; Hughes, G.; Massey, A. G. *Journal of Organometallic Chemistry*  
49  
50 **1976**, *112*, 9-19.  
51  
52
- 53 (34) Hara, H.; Watanabe, S.; Yamada, M.; Hoshino, O. *Journal of the Chemical Society,*  
54  
55 *Perkin Transactions 1: Organic and Bio-Organic Chemistry* **1996**, 741-745.  
56  
57  
58  
59  
60

- 1  
2  
3 (35) Ito, A.; Ono, Y.; Tanaka, K. *J. Org. Chem.* **1999**, *64*, 8236-8241.  
4  
5  
6 (36) Muci, A. R.; Buchwald, S. L. *Top. Curr. Chem.* **2002**, *219*, 131-209.  
7  
8 (37) Wolfe, J. P.; Wagaw, S.; Marcoux, J.; Buchwald, S. L. *Acc. Chem. Res.* **1998**, *31*,  
9  
10 805-818.  
11  
12 (38) Hartwig, J. F. *Angewandte Chemie, International Edition* **1998**, *37*, 2090-2093.  
13  
14 (39) Hartwig, J. F. *Acc. Chem. Res.* **1998**, *31*, 852-860.  
15  
16  
17 (40) Ley, S. V.; Thomas, A. W. *Angewandte Chemie, International Edition* **2003**, *42*,  
18  
19 5400-5449.  
20  
21  
22 (41) Bidman, T. A. *Russian Journal of General Chemistry (Translation of Zhurnal*  
23  
24 *Obshchei Khimii)* **2004**, *74*, 1433-1434.  
25  
26  
27 (42) Tietze, M.; Iglesias, A.; Merisor, E.; Conrad, J.; Klaiber, I.; Beifuss, U. *Org. Lett.*  
28  
29 **2005**, *7*, 1549-1552.  
30  
31  
32 (43) Black, D. S.; Rothnie, N. E. *Aust. J. Chem.* **1983**, *36*, 1141-1147.  
33  
34 (44) Wilshire, J. F. K. *Aust. J. Chem.* **1988**, *41*, 995-1001.  
35  
36 (45) Wilshire, J. F. K. *Australian Journal of Chemistry* **1988**, *41*, 995-1001.  
37  
38 (46) Sanz, R.; Fernandez, Y.; Castroviejo, M. P.; Perez, A.; Fananas, F. J. *J. Org. Chem.*  
39  
40 **2006**, *71*, 6291-6294.  
41  
42  
43 (47) Chong, H.; Garmestani, K.; Bryant, L. H., Jr.; Milenic, D. E.; Overstreet, T.; Birch,  
44  
45 N.; Le, T.; Brady, E. D.; Brechbiel, M. W. *J. Med. Chem.* **2006**, *49*, 2055-2062.  
46  
47  
48 (48) Levy, S. G.; Jacques, V.; Zhou, K. L.; Kalogeropoulos, S.; Schumacher, K.; Amedio,  
49  
50 J. C.; Scherer, J. E.; Witowski, S. R.; Lombardy, R.; Koppetsch, K. *Org. Process*  
51  
52 *Res. Dev.* **2009**, *13*, 535-542.  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 (49) Chong, H.; Ma, X.; Le, T.; Kwamena, B.; Milenic, D. E.; Brady, E. D.; Song, H. A.;  
4  
5 Brechbiel, M. W. *J. Med. Chem.* **2008**, *51*, 118.  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60