



8-2019

## Genome Sequence of Pseudomonas Phage UMP151, Isolated from the Female Bladder Microbiota

Catherine Putonti

Loyola University Chicago, [cputonti@luc.edu](mailto:cputonti@luc.edu)

Genevieve Johnson

Loyola University Chicago

Follow this and additional works at: [https://ecommons.luc.edu/bioinformatics\\_facpub](https://ecommons.luc.edu/bioinformatics_facpub)

 Part of the [Bioinformatics Commons](#)

---

### Recommended Citation

Putonti, Catherine and Johnson, Genevieve. Genome Sequence of Pseudomonas Phage UMP151, Isolated from the Female Bladder Microbiota. *Microbiology Resource Announcements*, , , 2019. Retrieved from Loyola eCommons, Bioinformatics Faculty Publications, <http://dx.doi.org/10.1128/MRA.00853-19>

This Article is brought to you for free and open access by the Faculty Publications and Other Works by Department at Loyola eCommons. It has been accepted for inclusion in Bioinformatics Faculty Publications by an authorized administrator of Loyola eCommons. For more information, please contact [ecommons@luc.edu](mailto:ecommons@luc.edu).



This work is licensed under a [Creative Commons Attribution 4.0 International License](#).  
Copyright © 2019 Johnson and Putonti.



# Genome Sequence of *Pseudomonas* Phage UMP151, Isolated from the Female Bladder Microbiota

Genevieve Johnson,<sup>a</sup>  Catherine Putonti<sup>a,b,c,d</sup>

<sup>a</sup>Bioinformatics Program, Loyola University Chicago, Chicago, Illinois, USA

<sup>b</sup>Department of Biology, Loyola University Chicago, Chicago, Illinois, USA

<sup>c</sup>Department of Computer Science, Loyola University Chicago, Chicago, Illinois, USA

<sup>d</sup>Department of Microbiology and Immunology, Loyola University Chicago, Maywood, Illinois, USA

**ABSTRACT** A temperate bacteriophage, designated UMP151, was isolated from a *Pseudomonas aeruginosa* strain from a catheterized urine sample of a woman with overactive bladder (OAB) symptoms. The 41,303-bp genome sequence of *Pseudomonas* phage UMP151 exhibits sequence similarity to prophage and lytic phage sequences isolated from other areas of the human body.

Prior studies of the female urinary microbiota have identified associations between bladder bacteria and urinary symptoms (1–6). Recent surveys have found that viruses, particularly bacteriophages (phages), far outnumber bacteria in the urinary tract (7) and that the majority of these bacterial members are lysogens, some harboring upwards of 10 intact prophages (8). In our ongoing effort to characterize the phage population of the urinary tract microbiota, we describe here the genome sequence of a new isolate, *Pseudomonas* phage UMP151, a temperate phage induced from a *Pseudomonas aeruginosa* strain isolated from the urine collected via transurethral catheterization from a woman with overactive bladder (OAB) symptoms. *P. aeruginosa* is not frequently found within the bladder microbiota of healthy women (9); rather, it is an opportunistic pathogen of the urinary tract typically associated with nosocomial urinary tract infections (10). Furthermore, no association between *P. aeruginosa* and OAB has been identified (9). *Pseudomonas* phage UMP151 is the fourth *Pseudomonas* phage isolated from the urinary tract to be described in the literature (11–13) and the first genome of a *Pseudomonas* phage from the bladder microbiota.

*P. aeruginosa* strain UMB0151 was isolated via culture from a prior study (1, 2, 6, 9) and stored at  $-80^{\circ}\text{C}$ . We streaked this bacterial strain on an LB agar plate and grew it overnight at  $37^{\circ}\text{C}$ . A single colony was selected to inoculate liquid LB medium and was grown overnight with shaking at  $37^{\circ}\text{C}$ . The overnight culture was centrifuged for 8 min at  $13,000 \times g$  and filtered through a  $0.2\text{-}\mu\text{m}$  cellulose acetate syringe filter, and then the filtrate was spotted ( $10\ \mu\text{l}$ ) onto a lawn of *P. aeruginosa* ATCC 15692. The plate was incubated overnight at  $37^{\circ}\text{C}$ . Plaques were found where the filtrate was spotted. Spots were harvested, suspended in LB medium, and filtered using a  $0.2\text{-}\mu\text{m}$  cellulose acetate syringe filter. A sample of  $300\ \mu\text{l}$  was pipetted and treated with 5 U of Optizyme DNase I (Fisher BioReagents) prior to DNA extraction using the Quick-DNA kit (Zymo) following the manufacturer's instructions. The Nextera XT DNA library preparation kit was used, and the library was sequenced on an Illumina MiSeq sequencer using the MiSeq reagent kit v2 (500 cycles), producing 278,404 paired-end 250-bp reads. Raw reads were first trimmed for quality using sickle (<https://github.com/najoshi/sickle>) and then assembled by SPAdes (v3.11.1) (14) using the “only assembler” option for  $k$  value of 55, 77, 99, and

**Citation** Johnson G, Putonti C. 2019. Genome sequence of *Pseudomonas* phage UMP151, isolated from the female bladder microbiota. Microbiol Resour Announc 8:e00853-19. <https://doi.org/10.1128/MRA.00853-19>.

**Editor** Simon Roux, DOE Joint Genome Institute

**Copyright** © 2019 Johnson and Putonti. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Catherine Putonti, [cputonti@luc.edu](mailto:cputonti@luc.edu).

**Received** 17 July 2019

**Accepted** 24 July 2019

**Published** 15 August 2019

127. In total, 4,828 assembled contigs were produced, ranging in size from 128 to 41,303 bp. Coverage was calculated using BBmap v38.47 (<http://sourceforge.net/projects/bbmap>). The largest contig, representative of the complete genome sequence of UMP151, had a significantly greater coverage (768 $\times$ ) than the other contigs assembled ( $\sim 1\times$ ). The genome termini were determined using PhageTerm v1.0.12 (15) via Galaxy (16), and the genome was annotated via RAST using the Classic RAST pipeline (17).

The complete genome for *Pseudomonas* phage UMP151 is 41,303 bp with a GC content of 63.2%. The genome encodes 56 genes. The complete genome sequence was queried against the NCBI nr/nt database via MegaBLAST, identifying 100% sequence identity and coverage with *P. aeruginosa* strain MRSN12280 (GenBank accession number CP028162), which was isolated from the sacrum of an individual. The best hit to an isolated phage was to the siphovirus *Pseudomonas* phage CF5 (GenBank accession number MK511057; query coverage, 77%; sequence identity, 97.48%), which was isolated from the lung of a cystic fibrosis patient (18).

**Data availability.** The draft whole-genome project for *Pseudomonas* phage UMP151 has been deposited at DDBJ/EMBL/GenBank under accession number MK934841. Raw sequence reads are deposited at DDBJ/EMBL/GenBank under accession number SRR9072121, which is part of BioProject number PRJNA494532.

## ACKNOWLEDGMENTS

We thank Alan J. Wolfe for the bacterial isolate and Roberto Limeira for whole-genome sequencing of the isolate at Loyola's Genomics Facility (Maywood, IL).

G.J.'s research was supported through the Loyola University Mulcahy Research Fellowship.

## REFERENCES

- Pearce MM, Hilt EE, Rosenfeld AB, Zilliox MJ, Thomas-White K, Fok C, Kliethermes S, Schreckenberger PC, Brubaker L, Gai X, Wolfe AJ. 2014. The female urinary microbiome: a comparison of women with and without urgency urinary incontinence. *mBio* 5:e01283-14. <https://doi.org/10.1128/mBio.01283-14>.
- Pearce MM, Zilliox MJ, Rosenfeld AB, Thomas-White KJ, Richter HE, Nager CW, Visco AG, Nygaard IE, Barber MD, Schaffer J, Moalli P, Sung VW, Smith AL, Rogers R, Nolen TL, Wallace D, Meikle SF, Gai X, Wolfe AJ, Brubaker L. 2015. The female urinary microbiome in urgency urinary incontinence. *Am J Obstet Gynecol* 213:347.e1–347.e11. <https://doi.org/10.1016/j.ajog.2015.07.009>.
- Fok CS, McKinley K, Mueller ER, Kenton K, Schreckenberger P, Wolfe A, Brubaker L. 2013. Day of surgery urine cultures identify urogynecologic patients at increased risk for postoperative urinary tract infection. *J Urol* 189:1721–1724. <https://doi.org/10.1016/j.juro.2012.11.167>.
- Karstens L, Asquith M, Davin S, Stauffer P, Fair D, Gregory WT, Rosenbaum JT, McWeeney SK, Nardos R. 2016. Does the urinary microbiome play a role in urgency urinary incontinence and its severity? *Front Cell Infect Microbiol* 6:78. <https://doi.org/10.3389/fcimb.2016.00078>.
- Nienhouse V, Gao X, Dong Q, Nelson DE, Toh E, McKinley K, Schreckenberger P, Shibata N, Fok CS, Mueller ER, Brubaker L, Wolfe AJ, Radek KA. 2014. Interplay between bladder microbiota and urinary antimicrobial peptides: mechanisms for human urinary tract infection risk and symptom severity. *PLoS One* 9:e114185. <https://doi.org/10.1371/journal.pone.0114185>.
- Thomas-White KJ, Hilt EE, Fok C, Pearce MM, Mueller ER, Kliethermes S, Jacobs K, Zilliox MJ, Brincat C, Price TK, Kuffel G, Schreckenberger P, Gai X, Brubaker L, Wolfe AJ. 2016. Incontinence medication response relates to the female urinary microbiota. *Int Urogynecol J* 27:723–733. <https://doi.org/10.1007/s00192-015-2847-x>.
- Garretto A, Miller-Ensminger T, Wolfe AJ, Putonti C. 2019. Bacteriophages of the lower urinary tract. *Nat Rev Urol* 16:422–432. <https://doi.org/10.1038/s41585-019-0192-4>.
- Miller-Ensminger T, Garretto A, Brenner J, Thomas-White K, Zambom A, Wolfe AJ, Putonti C. 2018. Bacteriophages of the urinary microbiome. *J Bacteriol* 200:e00738-17. <https://doi.org/10.1128/JB.00738-17>.
- Hilt EE, McKinley K, Pearce MM, Rosenfeld AB, Zilliox MJ, Mueller ER, Brubaker L, Gai X, Wolfe AJ, Schreckenberger PC. 2014. Urine is not sterile: use of enhanced urine culture techniques to detect resident bacterial flora in the adult female bladder. *J Clin Microbiol* 52:871–876. <https://doi.org/10.1128/JCM.02876-13>.
- Djordjevic Z, Folic MM, Zivic Z, Markovic V, Jankovic SM. 2013. Nosocomial urinary tract infections caused by *Pseudomonas aeruginosa* and *Acinetobacter* species: sensitivity to antibiotics and risk factors. *Am J Infect Control* 41:1182–1187. <https://doi.org/10.1016/j.ajic.2013.02.018>.
- Brown-Jaque M, Muniesa M, Navarro F. 2016. Bacteriophages in clinical samples can interfere with microbiological diagnostic tools. *Sci Rep* 6:33000. <https://doi.org/10.1038/srep33000>.
- Jalil MB, Al-Hmudi HA, Al-Asaad LA, Abdul-Hussein ZR. 2018. Isolation and characterization of bacteriophages against multiple drug resistant *Pseudomonas aeruginosa* with using the bacteriophage as a therapy in the mice model. *Int J Dev Res* 7:11519.
- Johnson G, Wolfe AJ, Putonti C. 2019. Characterization of the  $\phi$ CTX-like *Pseudomonas aeruginosa* phage Dobby isolated from the kidney stone microbiota. *Access Microbiol* 1. <https://doi.org/10.1099/acmi.0.000002>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Garneau JR, Depardieu F, Fortier L-C, Bikard D, Monot M. 2017. PhageTerm: a tool for fast and accurate determination of phage termini and packaging mechanism using next-generation sequencing data. *Sci Rep* 7:8292. <https://doi.org/10.1038/s41598-017-07910-5>.
- Afgan E, Baker D, Batut B, van den Beek M, Bouvier D, Čech M, Chilton J, Clements D, Coraor N, Grünig BA, Guerler A, Hillman-Jackson J, Hiltmann S, Jalili V, Rasche H, Soranzo N, Goecks J, Taylor J, Nekrutenko A, Blankenberg D. 2018. The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2018 update. *Nucleic Acids Res* 46:W537–W544. <https://doi.org/10.1093/nar/gky379>.
- Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K,

- Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <https://doi.org/10.1186/1471-2164-9-75>.
18. Tariq MA, Everest FLC, Cowley LA, Wright R, Holt GS, Ingram H, Duignan LAM, Nelson A, Lanyon CV, Perry A, Perry JD, Bourke S, Brockhurst MA, Bridge SH, De Soyza A, Smith DL. 2019. Temperate bacteriophages from chronic *Pseudomonas aeruginosa* lung infections show disease-specific changes in host range and modulate antimicrobial susceptibility. *mSystems* 4:e00191-18. <https://doi.org/10.1128/mSystems.00191-18>.