



6-4-2020

Draft Genome Sequence of Escherichia coli UMB9246, Isolated from the Bladder of a Woman with Recurrent Urinary Tract Infection

Mikalina Belmonte
Loyola University Chicago

Taylor Miller-Ensminger
Loyola University Chicago

Adelina Voukadinova
Loyola University Chicago

Alan J. Wolfe
Loyola University Chicago, awolfe@luc.edu

Catherine Putoni
Loyola University Chicago, cputonti@luc.edu

Follow this and additional works at: https://ecommons.luc.edu/bioinformatics_facpub

 Part of the [Bioinformatics Commons](#), and the [Biology Commons](#)

Recommended Citation

Belmonte, Mikalina; Miller-Ensminger, Taylor; Voukadinova, Adelina; Wolfe, Alan J.; and Putoni, Catherine. Draft Genome Sequence of Escherichia coli UMB9246, Isolated from the Bladder of a Woman with Recurrent Urinary Tract Infection. *Microbiology Resource Announcements*, 9, : , 2020. Retrieved from Loyola eCommons, Bioinformatics Faculty Publications, <http://dx.doi.org/10.1128/MRA.00437-20>

This Article is brought to you for free and open access by the Faculty Publications and Other Works by Department at Loyola eCommons. It has been accepted for inclusion in Bioinformatics Faculty Publications by an authorized administrator of Loyola eCommons. For more information, please contact ecommons@luc.edu.



This work is licensed under a [Creative Commons Attribution 4.0 License](#).
© Belmonte et al., 2020.



Draft Genome Sequence of *Escherichia coli* UMB9246, Isolated from the Bladder of a Woman with Recurrent Urinary Tract Infection

Mikalina Belmonte,^a Taylor Miller-Ensminger,^b Adelina Voukadinova,^b Alan J. Wolfe,^c  Catherine Putonti^{a,b,c,d}

^aDepartment of Biology, Loyola University Chicago, Chicago, Illinois, USA

^bBioinformatics Program, Loyola University Chicago, Chicago, Illinois, USA

^cDepartment of Microbiology and Immunology, Stritch School of Medicine, Loyola University Chicago, Maywood, Illinois, USA

^dDepartment of Computer Science, Loyola University Chicago, Chicago, Illinois, USA

ABSTRACT *Escherichia coli* is a Gram-negative, motile, rod-shaped bacterium that causes the majority of uncomplicated urinary tract infections (UTIs). Here, we report the draft genome of *E. coli* strain UMB9246, an isolate from a woman with recurrent UTI.

Pathogenic strains of *Escherichia coli* can cause intestinal, urinary tract, and internal infections, and *E. coli* isolates from multiple body sites have been sequenced (1). Specifically, uropathogenic *E. coli* (UPEC) strains are responsible for the majority of infrequent urinary tract infections (UTIs) in humans (2, 3). Recurrence of UTIs is frequent (4). Recurrent UTIs (rUTIs) are considerably more complicated and nuanced than infrequent UTIs. Here, we isolated *E. coli* strain UMB9246 from a catheterized urine sample obtained from a woman with an rUTI.

E. coli strain UMB9246 was collected as part of a prior institutional review board (IRB)-approved study (University of California, San Diego, IRB no. 170077AW) using the expanded quantitative urine culture (EQUC) protocol (5). Briefly, 100 μ l of urine was spread onto a 5% sheep blood agar plate and incubated overnight at 35°C with 5% CO₂ for 24 h. Each morphologically distinct colony was then further purified through successive plating. The genus and species for this isolate were determined via matrix-assisted laser desorption ionization–time of flight (MALDI-TOF) mass spectrometry as previously described (5). The isolate was then stored at –80°C until sequencing. From this freezer stock, the sample was streaked onto a Columbia nalidixic acid (CNA) agar plate and incubated at 35°C with 5% CO₂ for 24 h. A single colony was selected and incubated in LB broth at 37°C with shaking for 24 h. DNA was extracted using the DNeasy blood and tissue kit following the manufacturer’s protocol for Gram-positive bacteria with the following alterations. We used 230 μ l of lysis buffer (180 μ l of 20 mM Tris-Cl, 2 mM sodium EDTA, and 1.2% Triton X-100 and 50 μ l of lysozyme) in step 2, and in step 5 of the protocol, we incubated the sample at 56°C for 10 min. DNA was quantified using the Qubit fluorometer and sent to the Microbial Genome Sequencing Center (MiGS) at the University of Pittsburgh for sequencing. The DNA was enzymatically fragmented using an Illumina tagmentation enzyme, and indices were attached using PCR. The DNA was sequenced using an Illumina NextSeq 550 flow cell, producing 1,408,737 pairs of 150-bp reads. The raw reads were trimmed using Sickle v1.33 (<https://github.com/najoshi/sickle>) and assembled using SPAdes v3.13.0 with the “only-assembler” option for k values of 55, 77, 99, and 127 (6). The genome coverage is 71 \times and was calculated using BMap v38.47 (<https://sourceforge.net/projects/bbmap/>). The genome assembly was first annotated using PATRIC v3.6.3 (7). The NCBI Prokaryotic Genome Annotation Pipeline (PGAP) v4.11 (8) was used to annotate the publicly

Citation Belmonte M, Miller-Ensminger T, Voukadinova A, Wolfe AJ, Putonti C. 2020. Draft genome sequence of *Escherichia coli* UMB9246, isolated from the bladder of a woman with recurrent urinary tract infection. Microbiol Resour Announc 9:e00437-20. <https://doi.org/10.1128/MRA.00437-20>.

Editor David Rasko, University of Maryland School of Medicine

Copyright © 2020 Belmonte et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Catherine Putonti, cputonti@luc.edu.

Received 17 April 2020

Accepted 14 May 2020

Published 4 June 2020

available genome. The genome assembly was also examined using PHASTER (9) and CRISPRCasFinder (10) to identify prophage sequences and CRISPR spacer arrays, respectively. Default parameters were used for each software tool, unless previously stated.

The *E. coli* UMB9246 draft genome is 5,057,732 bp long in 90 contigs with an N_{50} score of 159,145 bp and a GC content of 50.8%. The PGAP annotation identified 4,691 protein-coding genes, 5 rRNA operons, and 77 tRNAs. PHASTER predicted 1 questionable, 2 incomplete, and 1 intact phage for this strain. CRISPRCasFinder found 2 CRISPR arrays. PATRIC identified in the genome assembly both virulence factors and antibiotic resistance genes, including those for tetracycline resistance. Further analysis of this strain and other isolates from individuals suffering from uncomplicated UTIs and rUTIs will give us insight into the mechanisms for recurrence.

Data availability. This whole-genome shotgun project has been deposited in GenBank under the accession no. [JAAUVS000000000](https://www.ncbi.nlm.nih.gov/nuclseq/JAAUVS000000000). The version described in this paper is the first version, JAAUVS010000000. The raw sequencing reads have been deposited in SRA under the accession no. [SRR11441020](https://www.ncbi.nlm.nih.gov/sra/SRR11441020).

ACKNOWLEDGMENTS

This work was conducted as part of the Bacterial Genomics course at Loyola University Chicago's Department of Biology. For prior patient recruitment, we acknowledge the Loyola Urinary Education and Research Collaborative (LUEREC) and the patients who provided the samples for this study.

REFERENCES

- Moriel DG, Tan L, Goh KGK, Phan M-D, Ipe DS, Lo AW, Peters KM, Ulett GC, Beatson SA, Schembri MA. 2016. A novel protective vaccine antigen from the core *Escherichia coli* genome. *mSphere* 1:e00326-16. <https://doi.org/10.1128/mSphere.00326-16>.
- Johnson JR. 1991. Virulence factors in *Escherichia coli* urinary tract infection. *Clin Microbiol Rev* 4:80–128. <https://doi.org/10.1128/cmr.4.1.80>.
- Terlizzi ME, Gribaudo G, Maffei ME. 2017. Uropathogenic *Escherichia coli* (UPEC) infections: virulence factors, bladder responses, antibiotic, and non-antibiotic antimicrobial strategies. *Front Microbiol* 8:1566. <https://doi.org/10.3389/fmicb.2017.01566>.
- Foxman B, Gillespie B, Koopman J, Zhang L, Palin K, Tallman P, Marsh JV, Spear S, Sobel JD, Marty MJ, Marrs CF. 2000. Risk factors for second urinary tract infection among college women. *Am J Epidemiol* 151:1194–1205. <https://doi.org/10.1093/oxfordjournals.aje.a010170>.
- Hilt EE, McKinley K, Pearce MM, Rosenfeld AB, Zilliox MJ, Mueller ER, Brubaker L, Gai X, Wolfe AJ, Schreckenberger PC. 2014. Urine is not sterile: use of enhanced urine culture techniques to detect resident bacterial flora in the adult female bladder. *J Clin Microbiol* 52:871–876. <https://doi.org/10.1128/JCM.02876-13>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Wattam AR, Davis JJ, Assaf R, Boisvert S, Brettin T, Bun C, Conrad N, Dietrich EM, Disz T, Gabbard JL, Gerdes S, Henry CS, Kenyon RW, Machi D, Mao C, Nordberg EK, Olsen GJ, Murphy-Olson DE, Olson R, Overbeek R, Parrello B, Pusch GD, Shukla M, Vonstein V, Warren A, Xia F, Yoo H, Stevens RL. 2017. Improvements to PATRIC, the all-bacterial Bioinformatics Database and Analysis Resource Center. *Nucleic Acids Res* 45:D535–D542. <https://doi.org/10.1093/nar/gkw1017>.
- Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
- Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, Wishart DS. 2016. PHASTER: a better, faster version of the PHAST phage search tool. *Nucleic Acids Res* 44:W16–W21. <https://doi.org/10.1093/nar/gkw387>.
- Couvin D, Bernheim A, Toffano-Nioche C, Touchon M, Michalik J, Néron B, Rocha EPC, Vergnaud G, Gautheret D, Pourcel C. 2018. CRISPRCas-Finder, an update of CRISPRFinder, includes a portable version, enhanced performance and integrates search for Cas proteins. *Nucleic Acids Res* 46:W246–W251. <https://doi.org/10.1093/nar/gky425>.