



Draft Genome Sequence of *Streptococcus anginosus* UMB7768, Isolated from a Woman with Recurrent UTI Symptoms

Sarah Miller,^a Taylor Miller-Ensminger,^a Adelina Voukadinova,^a Alan J. Wolfe,^b  Catherine Putonti^{a,b,c,d}

^aBioinformatics Program, Loyola University Chicago, Chicago, Illinois, USA

^bDepartment of Microbiology and Immunology, Stritch School of Medicine, Loyola University Chicago, Maywood, Illinois, USA

^cDepartment of Biology, Loyola University Chicago, Chicago, Illinois, USA

^dDepartment of Computer Science, Loyola University Chicago, Chicago, Illinois, USA

ABSTRACT *Streptococcus anginosus* recently was implicated as a pathogen involved in urinary tract infections. A strain of *S. anginosus* was isolated from the female urogenital tract. Here, we present the draft genome sequence of *S. anginosus* strain UMB7768.

Streptococcus anginosus is a catalase-negative Gram-positive bacterium normally associated with a beneficial role in the human gastrointestinal and oropharynx microbiota (1, 2). However, recent studies have shown that *S. anginosus* is a pathogenic contributor to several abscess-forming infections and potentially plays a role in cystic fibrosis (3, 4). Furthermore, *S. anginosus* is the predominant member of the *S. milleri* group (which also includes *S. constellatus* and *S. intermedius*) found in the urogenital tract (5). A prior study found that strains of the bladder and vaginal communities were highly similar, suggesting microbial sharing between these two microbiota (6). *S. anginosus* is considered an emerging uropathogen; it has been associated with urgency urinary incontinence (7) and urinary tract infections (UTIs) (8, 9). Here, we present the draft genome sequence of *S. anginosus* UMB7768 from the urinary microbiota of a female with recurrent UTI.

A voided urine specimen was collected from a patient at the Women's Pelvic Medicine Center at the University of California, San Diego, as part of a previous institutional review board (IRB)-approved study (University of California, San Diego, IRB no. 170077AW). *S. anginosus* UMB7768 was isolated from this sample using the expanded quantitative urinary culture (EQUC) protocol (10). The genus and species of this isolate were determined with matrix-assisted laser desorption ionization–time of flight (MALDI-TOF) mass spectrometry (10) and then stored at -80°C . Following freezing, the isolate was streaked onto a Columbia nalidixic acid (CNA) agar plate and incubated at 35°C with 5% CO_2 for 24 h. A single colony was selected from the plate and inoculated in blood heart infusion (BHI) broth at 37°C with shaking for 24 h. DNA was extracted using the Qiagen DNeasy blood and tissue kit following the manufacturer's protocol for Gram-positive bacteria (with minor modification) and quantified using a Qubit fluorometer. The extracted DNA was sent to the Microbial Genomic Sequencing Center (MiGS) at the University of Pittsburgh for sequencing. There, the DNA was enzymatically fragmented using an Illumina tagmentation enzyme, and then indices were attached using PCR. The DNA library was sequenced using an Illumina NextSeq 550 platform, producing 1,513,634 pairs of 150-bp reads. Raw reads were trimmed using Sickle v1.33 (<https://github.com/najoshi/sickle>) and assembled using SPAdes v3.13 with the “only assembler” option for k values of 55, 77, 99, and 127 (11). The genome coverage was calculated using BBMap v38.47 (<https://sourceforge.net/projects/bbmap/>). PATRIC v3.6.3 (12) was used to annotate the genome in-house; the publicly available genome

Citation Miller S, Miller-Ensminger T, Voukadinova A, Wolfe AJ, Putonti C. 2020. Draft genome sequence of *Streptococcus anginosus* UMB7768, isolated from a woman with recurrent UTI symptoms. Microbiol Resour Announc 9:e00418-20. <https://doi.org/10.1128/MRA.00418-20>.

Editor Julie C. Dunning Hotopp, University of Maryland School of Medicine

Copyright © 2020 Miller et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Catherine Putonti, cputonti@luc.edu.

Received 15 April 2020

Accepted 28 April 2020

Published 21 May 2020

sequence was annotated using the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) v4.11 (13). Unless otherwise noted, default parameters were used for all software tools listed.

The *S. anginosus* UMB7768 draft genome sequence is 2,158,792 bp long, assembled in 47 contigs with an N_{50} value of 121,537 bp and a GC content of 38.77%. The genome's coverage is 182 \times . The PGAP annotation found 2,168 coding DNA sequences (CDSs), 2,000 with protein prediction, 45 tRNAs, and 5 complete rRNA sequences (3 5S, 1 16S, and 1 23S). PATRIC identified numerous virulence factors and genes associated with antibiotic resistance. Further investigation of the latter using the ResFinder v3.2 (14) Web server identified resistances to macrolides and tetracycline, which have been documented as common within the *S. milleri* group (4). Future examination of the virulence factors identified will expand our understanding of *S. anginosus* in the urogenital tract and its role in UTI development.

Data availability. This whole-genome shotgun project has been deposited in GenBank under the accession no. [JAAUWB000000000](https://www.ncbi.nlm.nih.gov/nuclseq/JAAUWB000000000). The version described in this paper is the first version, JAAUWB010000000. The raw sequencing reads have been deposited in the SRA under the accession no. [SRR11441032](https://www.ncbi.nlm.nih.gov/sra/SRR11441032).

ACKNOWLEDGMENTS

This work was conducted as part of Loyola University Chicago's Department of Biology Bacterial Genomics course. For prior patient recruitment, we acknowledge the Loyola Urinary Education and Research Collaborative (LUEREC) and the patients who provided the samples for this study.

REFERENCES

1. Spellerberg B, Brandt C. 2015. Streptococcus, p 383–402. In Jorgenson JH, Pfaller MA, Carroll KC, Funke G, Landry ML, Richter SS, Warnock DW (ed), *Manual of clinical microbiology*, 11th ed. ASM Press, Washington, DC.
2. Whiley RA, Beighton D, Winstanley TG, Fraser HY, Hardie JM. 1992. Streptococcus intermedius, Streptococcus constellatus, and Streptococcus anginosus (the Streptococcus milleri group): association with different body sites and clinical infections. *J Clin Microbiol* 30:243–244. <https://doi.org/10.1128/JCM.30.1.243-244.1992>.
3. Asam D, Spellerberg B. 2014. Molecular pathogenicity of Streptococcus anginosus. *Mol Oral Microbiol* 29:145–155. <https://doi.org/10.1111/omi.12056>.
4. Grinwis ME, Sibley CD, Parkins MD, Eshaghurshan CS, Rabin HR, Surette MG. 2010. Macrolide and clindamycin resistance in Streptococcus milleri group isolates from the airways of cystic fibrosis patients. *Antimicrob Agents Chemother* 54:2823–2829. <https://doi.org/10.1128/AAC.01845-09>.
5. Jakubovics NS, Yassin SA, Rikard AH. 2014. Community interactions of oral streptococci. *Adv Appl Microbiol* 87:43–110. <https://doi.org/10.1016/B978-0-12-800261-2.00002-5>.
6. Thomas-White K, Forster SC, Kumar N, Van Kuiken M, Putonti C, Stares MD, Hilt EE, Price TK, Wolfe AJ, Lawley TD. 2018. Culturing of female bladder bacteria reveals an interconnected urogenital microbiota. *Nat Commun* 9:1557. <https://doi.org/10.1038/s41467-018-03968-5>.
7. Pearce MM, Hilt EE, Rosenfeld AB, Zilliox MJ, Thomas-White K, Fok C, Kliethermes S, Schreckenberger PC, Brubaker L, Gai X, Wolfe AJ. 2014. The female urinary microbiome: a comparison of women with and without urgency urinary incontinence. *mBio* 5:e01283-14. <https://doi.org/10.1128/mBio.01283-14>.
8. Siegman-Igra Y, Azmon Y, Schwartz D. 2012. Milleri group streptococcus: a stepchild in the viridans family. *Eur J Clin Microbiol Infect Dis* 31:2453–2459. <https://doi.org/10.1007/s10096-012-1589-7>.
9. Furuichi M, Horikoshi Y. 2018. Sites of infection associated with Streptococcus anginosus group among children. *J Infect Chemother* 24:99–102. <https://doi.org/10.1016/j.jiac.2017.09.011>.
10. Hilt EE, McKinley K, Pearce MM, Rosenfeld AB, Zilliox MJ, Mueller ER, Brubaker L, Gai X, Wolfe AJ, Schreckenberger PC. 2014. Urine is not sterile: use of enhanced urine culture techniques to detect resident bacterial flora in the adult female bladder. *J Clin Microbiol* 52:871–876. <https://doi.org/10.1128/JCM.02876-13>.
11. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
12. Brettin T, Davis JJ, Disz T, Edwards RA, Gerdes S, Olsen GJ, Olson R, Overbeek R, Parrello B, Pusch GD, Shukla M, Thomason JA, 3rd, Stevens R, Vonstein V, Wattam AR, Xia F. 2015. RASTtk: a modular and extensible implementation of the RAST algorithm for building custom annotation pipelines and annotating batches of genomes. *Sci Rep* 5:8365. <https://doi.org/10.1038/srep08365>.
13. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
14. Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, Aarestrup FM, Larsen MV. 2012. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother* 67:2640–2644. <https://doi.org/10.1093/jac/dks261>.