



5-21-2020

Draft Genome Sequences of Three *Enterococcus casseliflavus* Strains Isolated from the Urine of Healthy Bovine Heifers (Gyr Breed)

Silvia Giannattasio-Ferraz
Loyola University Chicago

Laura Maskeri
Loyola University Chicago

André P. Oliveira
Loyola University Chicago

Edel F. Barbosa-Stancioli
Loyola University Chicago

Catherine Putonti
Loyola University Chicago, cputonti@luc.edu

Follow this and additional works at: https://ecommons.luc.edu/bioinformatics_facpub

 Part of the [Bioinformatics Commons](#), and the [Biology Commons](#)

Recommended Citation

Giannattasio-Ferraz, Silvia; Maskeri, Laura; Oliveira, André P.; Barbosa-Stancioli, Edel F.; and Putonti, Catherine. Draft Genome Sequences of Three *Enterococcus casseliflavus* Strains Isolated from the Urine of Healthy Bovine Heifers (Gyr Breed). *Microbiology Resource Announcements*, 9, : , 2020. Retrieved from Loyola eCommons, Bioinformatics Faculty Publications, <http://dx.doi.org/10.1128/MRA.00386-20>

This Article is brought to you for free and open access by the Faculty Publications and Other Works by Department at Loyola eCommons. It has been accepted for inclusion in Bioinformatics Faculty Publications by an authorized administrator of Loyola eCommons. For more information, please contact ecommons@luc.edu.



This work is licensed under a [Creative Commons Attribution 4.0 International License](#).

© Giannattasio-Ferraz et al., 2020



Draft Genome Sequences of Three *Enterococcus casseliflavus* Strains Isolated from the Urine of Healthy Bovine Heifers (Gyr Breed)

Silvia Giannattasio-Ferraz,^a Laura Maskeri,^b André P. Oliveira,^c Edel F. Barbosa-Stancioli,^a  Catherine Putonti^{b,d,e,f}

^aDepartamento de Microbiologia, Instituto de Ciências Biológicas, Universidade Federal Minas Gerais, Belo Horizonte, MG, Brazil

^bBioinformatics Program, Loyola University Chicago, Chicago, Illinois, USA

^cEmpresa de Pesquisa Agropecuária de Minas Gerais, Uberaba, MG, Brazil

^dDepartment of Biology, Loyola University Chicago, Chicago, Illinois, USA

^eDepartment of Computer Science, Loyola University Chicago, Chicago, Illinois, USA

^fDepartment of Microbiology and Immunology, Stritch School of Medicine, Loyola University Chicago, Maywood, Illinois, USA

ABSTRACT *Enterococcus casseliflavus* is a commensal bacterium present in the intestinal microbiota of different animals. Previous studies have found that strains isolated from livestock are often resistant to many different antibiotics. Here, we present three *E. casseliflavus* strains, UFMG-H7, UFMG-H8, and UFMG-H9, isolated from urine collected from healthy dairy heifers in Brazil.

Enterococcus casseliflavus is a commensal of the intestinal tract of livestock animals, such as cattle, horses, and sheep (1, 2). Antibiotic resistance is prevalent within this species, with isolates from livestock animals being reported as resistant to vancomycin, tetracycline, erythromycin, ampicillin, and gentamicin (3–5). Here, we report the draft genome sequences of three strains of *E. casseliflavus*, strains UFMG-H7, UFMG-H8, and UFMG-H9, which were isolated from urine collected from healthy dairy cattle in Brazil.

Sample collection took place at the Agricultural Research Company of Minas Gerais State in May 2019 and was previously approved by the Ethics Committee in Animal Experimentation of the Universidade Federal de Minas Gerais, Brazil (CEUA/UFMG approval number 40/2019). All of the assays performed were in accordance with relevant guidelines. The heifers sampled were from a herd composed of pure-by-origin Gyr cattle. For sampling, the vulva of the heifer was washed with distilled water and soap. Midstream urine was collected using a 50-ml sterile tube. The material was kept at –20°C for no more than 48 h, when it was brought back to the laboratory for processing. Aliquots of 2 ml were centrifuged, and the supernatant was plated on lysogeny broth (LB) agar plates. Plates were incubated overnight at 37°C, and then single colonies were grown in LB medium overnight at 37°C. This process of plating and liquid culture growth was repeated three times until pure colonies were obtained. A single colony was inoculated in LB liquid medium and grown overnight at 37°C under agitation. The DNA was extracted using the Qiagen DNeasy UltraClean microbial kit (Qiagen, Hilden, Germany) and quantified using a Qubit fluorometer. The 16S rRNA gene sequences were amplified using the primers 63f and 1387r (6) and sequenced by GENEWIZ (South Plainfield, NJ, USA) via Sanger sequencing using each primer. The 16S rRNA gene sequences produced were then queried against the NCBI 16S rRNA sequence database to ascertain the species of the isolates. Purified DNA was subjected to whole-genome sequencing at the Microbial Genomic Sequencing Center (MiGS) at the University of Pittsburgh. DNA libraries were prepared by the MiGS as follows: the DNA was fragmented using a tagmentation enzyme (Illumina), and indices were attached by PCR. Genome sequencing was conducted using the NextSeq 550 platform.

Citation Giannattasio-Ferraz S, Maskeri L, Oliveira AP, Barbosa-Stancioli EF, Putonti C. 2020. Draft genome sequences of three *Enterococcus casseliflavus* strains isolated from the urine of healthy bovine heifers (Gyr breed). Microbiol Resour Announc 9:e00386-20. <https://doi.org/10.1128/MRA.00386-20>.

Editor Christina A. Cuomo, Broad Institute

Copyright © 2020 Giannattasio-Ferraz et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Catherine Putonti, cputonti@luc.edu.

Received 12 April 2020

Accepted 27 April 2020

Published 21 May 2020

TABLE 1 Genome statistics for *E. casseliflavus* strains

Parameter	Data for strain:		
	UFMG-H7	UFMG-H8	UFMG-H9
Length (bp)	3,545,372	3,635,178	3,622,925
No. of contigs	19	26	26
Genome coverage (×)	47	111	50
N_{50} (bp)	282,784	503,206	447,071
No. of raw read pairs	1,423,449	1,464,825	1,405,112
GC content (%)	45.32	41.03	45.69
No. of coding genes	3,238	3,347	3,315
No. of tRNAs	53	53	53
No. of plasmids	1	1	0

Sequencing produced 2×150 -bp reads. These reads were trimmed using Sickle v1.33 (<https://github.com/najoshi/sickle>) and assembled using SPAdes v3.13.0 with the only-assembler option for k values of 55, 77, 99, and 127 (7). After assembly, the genome coverage was calculated using BMap v38.47 (<https://sourceforge.net/projects/bbmap>), and the genomes were annotated by the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) v4.11 (8). The genomes were examined for plasmids using PlasmidFinder v2.0 (9) and for antibiotic resistance genes using ResFinder v3.2 (10). Default parameters were used for all software tools unless otherwise noted.

Table 1 lists the genome statistics for the three *E. casseliflavus* bovine urinary isolates. Both UFMG-H7 and UFMG-H8 were found to contain an Inc18 plasmid (67.7 kbp and 36.7 kbp long, respectively). UFMG-H9 was not found to contain a plasmid. All three strains were found to contain vancomycin resistance genes. These genes were not carried by plasmids; rather, they were contained in the chromosome. No other antibiotic resistance genes were detected. Further analyses for better understanding of the circulation and role of these bacteria within the urinary tract are relevant to improve our knowledge regarding the bovine microbiota.

Data availability. These whole-genome sequencing projects are available in GenBank under the accession numbers [JAAVMT000000000](https://ncbi.nlm.nih.gov/nucl/JAAVMT000000000) (UFMG-H7), [JAAVMP000000000](https://ncbi.nlm.nih.gov/nucl/JAAVMP000000000) (UFMG-H8), and [JAAVMR000000000](https://ncbi.nlm.nih.gov/nucl/JAAVMR000000000) (UFMG-H9). Raw data are available under SRA accession numbers [SRR11455848](https://ncbi.nlm.nih.gov/sra/SRR11455848) (UFMG-H7), [SRR11455647](https://ncbi.nlm.nih.gov/sra/SRR11455647) (UFMG-H8), and [SRR11455846](https://ncbi.nlm.nih.gov/sra/SRR11455846) (UFMG-H9). These genome sequences are part of BioProject accession number [PRJNA615899](https://ncbi.nlm.nih.gov/bioproject/PRJNA615899).

REFERENCES

- Graves A, Weaver RW, Entry J. 2009. Characterization of enterococci populations in livestock manure using BIOLOG. *Microbiol Res* 164: 260–266. <https://doi.org/10.1016/j.micres.2006.11.017>.
- Beukers AG, Zaheer R, Goji N, Amoako KK, Chaves AV, Ward MP, McAllister TA. 2017. Comparative genomics of *Enterococcus* spp. isolated from bovine feces. *BMC Microbiol* 17:52. <https://doi.org/10.1186/s12866-017-0962-1>.
- Ngbede EO, Raji MA, Kwanashie CN, Kwaga J. 2017. Antimicrobial resistance and virulence profile of enterococci isolated from poultry and cattle sources in Nigeria. *Trop Anim Health Prod* 49:451–458. <https://doi.org/10.1007/s11250-016-1212-5>.
- Iweriebor BC, Obi LC, Okoh AI. 2016. Macrolide, glycopeptide resistance and virulence genes in *Enterococcus* species isolates from dairy cattle. *J Med Microbiol* 65:641–648. <https://doi.org/10.1099/jmm.0.000275>.
- de Jong A, Simjee S, Rose M, Moyaert H, El Garch F, Youala M. 2019. Antimicrobial resistance monitoring in commensal enterococci from healthy cattle, pigs and chickens across Europe during 2004–14 (EASSA Study). *J Antimicrob Chemother* 74:921–930. <https://doi.org/10.1093/jac/dky537>.
- Marchesi JR, Sato T, Weightman AJ, Martin TA, Fry JC, Hiom SJ, Dymock D, Wade WG. 1998. Design and evaluation of useful bacterium-specific PCR primers that amplify genes coding for bacterial 16S rRNA. *Appl Environ Microbiol* 64:795–799. <https://doi.org/10.1128/AEM.64.2.795-799.1998>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
- Carattoli A, Zankari E, García-Fernández A, Voldby Larsen M, Lund O, Villa L, Møller Aarestrup F, Hasman H. 2014. In silico detection and typing of plasmids using PlasmidFinder and plasmid multilocus sequence typing. *Antimicrob Agents Chemother* 58:3895–3903. <https://doi.org/10.1128/AAC.02412-14>.
- Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, Aarestrup FM, Larsen MV. 2012. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother* 67:2640–2644. <https://doi.org/10.1093/jac/dks261>.