



5-14-2020

Draft Genome Sequence of Antibiotic-Resistant *Enterococcus faecalis* Strain UMB0843, Isolated from the Female Urinary Tract

Natalia Purta
Loyola University Chicago

Taylor Miller-Ensminger
Loyola University Chicago

Adelina Voukadinova
Loyola University Chicago

Alan J. Wolfe
Loyola University Chicago, awolfe@luc.edu

Catherine Putoni
Loyola University Chicago, cputonti@luc.edu

Follow this and additional works at: https://ecommons.luc.edu/bioinformatics_facpub



Part of the [Bioinformatics Commons](#), and the [Biology Commons](#)

Recommended Citation

Purta, Natalia; Miller-Ensminger, Taylor; Voukadinova, Adelina; Wolfe, Alan J.; and Putoni, Catherine. Draft Genome Sequence of Antibiotic-Resistant *Enterococcus faecalis* Strain UMB0843, Isolated from the Female Urinary Tract. *Microbiology Resource Announcements*, 9, : , 2020. Retrieved from Loyola eCommons, Bioinformatics Faculty Publications, <http://dx.doi.org/10.1128/MRA.00407-20>

This Article is brought to you for free and open access by the Faculty Publications and Other Works by Department at Loyola eCommons. It has been accepted for inclusion in Bioinformatics Faculty Publications by an authorized administrator of Loyola eCommons. For more information, please contact ecommons@luc.edu.



This work is licensed under a [Creative Commons Attribution 4.0 License](#).

© Purta et al., 2020



Draft Genome Sequence of Antibiotic-Resistant *Enterococcus faecalis* Strain UMB0843, Isolated from the Female Urinary Tract

Natalia Purta,^a Taylor Miller-Ensminger,^b Adelina Voukadinova,^b Alan J. Wolfe,^c  Catherine Putonti^{b,c,d,e}

^aNeuroscience Program, Loyola University Chicago, Chicago, Illinois, USA

^bBioinformatics Program, Loyola University Chicago, Chicago, Illinois, USA

^cDepartment of Microbiology and Immunology, Stritch School of Medicine, Loyola University Chicago, Maywood, Illinois, USA

^dDepartment of Biology, Loyola University Chicago, Chicago, Illinois, USA

^eDepartment of Computer Science, Loyola University Chicago, Chicago, Illinois, USA

ABSTRACT Here, we introduce the 2.8-Mbp draft genome of *Enterococcus faecalis* strain UMB0843, isolated from the female urinary tract. *E. faecalis* is a leading cause of nosocomial infections, and many strains are often resistant to multiple antibiotics. We focus our genome analysis on the multiple genes involved in antibiotic resistance in this strain.

Enterococcus faecalis lives in the gastrointestinal tract of many organisms (1) and has recently emerged as a leading cause of nosocomial infections due to its increased resistance to many antibiotics (2, 3). *E. faecalis* is often a primary cause of surgical infections, infections within the bloodstream, and urinary tract infections (UTIs) (3, 4). Recently, we isolated *E. faecalis* strain UMB0843 from a urine sample obtained from a pregnant female. Here, we present the draft genome sequence of this isolate and the antibiotic resistance genes found within the genome.

E. faecalis UMB0843 was collected as part of a previous institutional review board (IRB)-approved study (5) and cultured using the expanded quantitative urine culture (EQUC) protocol (6). To determine the genus and species of each isolate, matrix-assisted laser desorption ionization–time of flight (MALDI-TOF) mass spectrometry was used, following a protocol detailed previously (6), prior to storage of the isolates at -80°C . From the freezer stocks, *E. faecalis* was streaked onto a Columbia nalidixic acid (CNA) agar plate and incubated at 35°C in 5% CO_2 for 24 h. A single colony was selected, inoculated in brain heart infusion (BHI) broth (premixed BBL brain heart infusion; BD), and incubated under the same conditions described above. DNA was extracted using the Qiagen DNeasy blood and tissue kit and quantified using the fluorescence-based Qubit. The Gram-positive extraction protocol was followed with the following exception: 230 μl of lysis buffer was used (180 μl of 20 mM Tris-Cl, 2 mM sodium EDTA, and 1.2% Triton X-100 and 50 μl of lysozyme). The extracted DNA was sent to the Microbial Genome Sequencing Center (MiGS) at the University of Pittsburgh for sequencing. The DNA was first enzymatically fragmented using the Illumina tagmentation enzyme, and then indices were attached using PCR. Sequencing was performed using an Illumina NextSeq 550 flow cell, and 1,590,783 pairs of 150-bp reads were generated. The raw reads were trimmed using Sickle v1.33 (<https://github.com/najoshi/sickle>) and assembled using SPAdes v3.13.0 with the “only-assembler” option for k values of 55, 77, 99, and 127 (7). BMap v38.47 (<https://sourceforge.net/projects/bbmap/>) was used to calculate the genome coverage. Genome sequences were annotated using PATRIC v3.6.3 (8), while the publicly available genome was annotated using the NCBI Prokary-

Citation Purta N, Miller-Ensminger T, Voukadinova A, Wolfe AJ, Putonti C. 2020. Draft genome sequence of antibiotic-resistant *Enterococcus faecalis* strain UMB0843, isolated from the female urinary tract. Microbiol Resour Announc 9:e00407-20. <https://doi.org/10.1128/MRA.00407-20>.

Editor John J. Dennehy, Queens College

Copyright © 2020 Purta et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Catherine Putonti, cputonti@luc.edu.

Received 13 April 2020

Accepted 27 April 2020

Published 14 May 2020

otic Genome Annotation Pipeline (PGAP) v4.11 (9). Unless stated otherwise, default parameters were used for all software.

The *E. faecalis* UMB0843 draft genome is 2,805,168 bp long, assembled into 20 contigs with a genome coverage of 147 \times and an N_{50} score of 284,798 bp. The genome has a GC content of 38%, similar to that of other genomes of the species. PGAP identified 2,611 protein-coding regions. Both annotation tools found 51 tRNAs and 5 complete rRNA gene sequences (3 5S, 1 16S, and 1 23S). While resistance of this strain was not experimentally tested, PATRIC reported 39 genes associated with antibiotic resistance. Upon further investigation using ResFinder v3.2 (10), only antibiotic resistance to macrolides was detected [resistance gene, *Isa(A)*]. Our analysis suggests that this strain is susceptible to vancomycin, commonly used as the last line of defense. Vancomycin-resistant strains of *E. faecalis* emerged in the United States first in 1989 (11) and have become a significant concern for UTI treatment (12).

Data availability. This whole-genome shotgun project has been deposited in DDBJ/ENA/GenBank under the accession no. [JAAUWL000000000](https://doi.org/10.1093/nar/gkw1017). The version described in this paper is the first version, JAAUWL010000000. The raw sequencing reads have been deposited in the SRA under the accession no. [SRR11441019](https://doi.org/10.1093/nar/gkw1017).

ACKNOWLEDGMENTS

This work was conducted as part of the Bacterial Genomics course at Loyola University Chicago's Department of Biology. For prior patient recruitment, we acknowledge the Loyola Urinary Education and Research Collaborative (LUEREC) and the patients who provided the samples for this study.

REFERENCES

1. Lawley TD, Walker AW. 2013. Intestinal colonization resistance. *Immunology* 138:1–11. <https://doi.org/10.1111/j.1365-2567.2012.03616.x>.
2. Gilmore MS, Lebreton F, van Schaik W. 2013. Genomic transition of enterococci from gut commensals to leading causes of multidrug-resistant hospital infection in the antibiotic era. *Curr Opin Microbiol* 16:10–16. <https://doi.org/10.1016/j.mib.2013.01.006>.
3. Jett BD, Huycke MM, Gilmore MS. 1994. Virulence of enterococci. *Clin Microbiol Rev* 7:462–478. <https://doi.org/10.1128/cmr.7.4.462>.
4. Hidron AI, Edwards JR, Patel J, Horan TC, Sievert DM, Pollock DA, Fridkin SK, National Healthcare Safety Network Team, Participating National Healthcare Safety Network Facilities. 2008. Antimicrobial-resistant pathogens associated with healthcare-associated infections: annual summary of data reported to the National Healthcare Safety Network at the Centers for Disease Control and Prevention, 2006–2007. *Infect Control Hosp Epidemiol* 29:996–1011. <https://doi.org/10.1086/591861>.
5. Jacobs KM, Thomas-White KJ, Hilt EE, Wolfe AJ, Waters TP. 2017. Microorganisms identified in the maternal bladder: discovery of the maternal bladder microbiota. *AJP Rep* 7:e188–e196. <https://doi.org/10.1055/s-0037-1606860>.
6. Hilt EE, McKinley K, Pearce MM, Rosenfeld AB, Zilliox MJ, Mueller ER, Brubaker L, Gai X, Wolfe AJ, Schreckenberger PC. 2014. Urine is not sterile: use of enhanced urine culture techniques to detect resident bacterial flora in the adult female bladder. *J Clin Microbiol* 52:871–876. <https://doi.org/10.1128/JCM.02876-13>.
7. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyskhin AV, Sirotkin AV, Yahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
8. Wattam AR, Davis JJ, Assaf R, Boisvert S, Brettin T, Bun C, Conrad N, Dietrich EM, Disz T, Gabbard JL, Gerdes S, Henry CS, Kenyon RW, Machi D, Mao C, Nordberg EK, Olsen GJ, Murphy-Olson DE, Olson R, Overbeek R, Parrello B, Pusch GD, Shukla M, Vonstein V, Warren A, Xia F, Yoo H, Stevens RL. 2017. Improvements to PATRIC, the all-bacterial Bioinformatics Database and Analysis Resource Center. *Nucleic Acids Res* 45:D535–D542. <https://doi.org/10.1093/nar/gkw1017>.
9. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
10. Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, Aarestrup FM, Larsen MV. 2012. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother* 67:2640–2644. <https://doi.org/10.1093/jac/dks261>.
11. Sahm DF, Kissinger J, Gilmore MS, Murray PR, Mulder R, Solliday J, Clarke B. 1989. In vitro susceptibility studies of vancomycin-resistant *Enterococcus faecalis*. *Antimicrob Agents Chemother* 33:1588–1591. <https://doi.org/10.1128/aac.33.9.1588>.
12. Shrestha LB, Baral R, Khanal B. 2019. Comparative study of antimicrobial resistance and biofilm formation among Gram-positive uropathogens isolated from community-acquired urinary tract infections and catheter-associated urinary tract infections. *Infect Drug Resist* 12:957–963. <https://doi.org/10.2147/IDR.S200988>.