



5-14-2020

Draft Genome Sequence of *Corynebacterium coyleae* UMB8490, Isolated from the Female Urinary Tract

Abbas Ali Khan
Loyola University Chicago

Taylor Miller-Ensminger
Loyola University Chicago

Adelina Voukadinova
Loyola University Chicago

Alan J. Wolfe
Loyola University Chicago, awolfe@luc.edu

Catherine Putonti
Loyola University Chicago, cputonti@luc.edu

Follow this and additional works at: https://ecommons.luc.edu/bioinformatics_facpub



Part of the [Bioinformatics Commons](#), and the [Biology Commons](#)

Recommended Citation

Ali Khan, Abbas; Miller-Ensminger, Taylor; Voukadinova, Adelina; Wolfe, Alan J.; and Putonti, Catherine. Draft Genome Sequence of *Corynebacterium coyleae* UMB8490, Isolated from the Female Urinary Tract. *Microbiology Resource Announcements*, 9, : , 2020. Retrieved from Loyola eCommons, Bioinformatics Faculty Publications, <http://dx.doi.org/10.1128/MRA.00393-20>

This Article is brought to you for free and open access by the Faculty Publications and Other Works by Department at Loyola eCommons. It has been accepted for inclusion in Bioinformatics Faculty Publications by an authorized administrator of Loyola eCommons. For more information, please contact ecommons@luc.edu.



This work is licensed under a [Creative Commons Attribution 4.0 International License](#).

© Ali Khan et al., 2020



Draft Genome Sequence of *Corynebacterium coyleae* UMB8490, Isolated from the Female Urinary Tract

Abbas Ali Khan,^a Taylor Miller-Ensminger,^b Adelina Voukadinova,^b Alan J. Wolfe,^c  Catherine Putonti^{a,b,c,d}

^aDepartment of Biology, Loyola University Chicago, Chicago, Illinois, USA

^bBioinformatics Program, Loyola University Chicago, Chicago, Illinois, USA

^cDepartment of Microbiology and Immunology, Stritch School of Medicine, Loyola University Chicago, Maywood, Illinois, USA

^dDepartment of Computer Science, Loyola University Chicago, Chicago, Illinois, USA

ABSTRACT Here, we present the draft genome sequence for *Corynebacterium coyleae* UMB8490, isolated from the catheterized urine of a female with overactive bladder symptoms. The size of the genome is 2,483,223 bp assembled in 62 contigs with an observed GC content of 61.42%.

Corynebacterium is a genus of rod-shaped, Gram-positive, facultative anaerobes (1). The species *Corynebacterium coyleae* has been associated with urinary tract infections (UTIs), sometimes being the single culture-positive species detected (2). *C. coyleae* has also been associated with urgency urinary incontinence (UUI) and overactive bladder (OAB) (3, 4). *C. coyleae* has also been found in the urine cultures of healthy individuals, but generally in lower abundances, indicating that it may be part of the normal microbiota of the bladder (2, 5). While there is scarce information on the pathogenic potential of this species, it is considered to be an emerging pathogen involved in complicated UTIs as well as nosocomial infections (2, 6).

C. coyleae UMB8490 was isolated from a catheterized urine sample obtained from a female patient with OAB as part of a prior institutional review board (IRB)-approved study (University of California, San Diego, IRB no. 170077AW) using the expanded quantitative urinary culture (EQUC) protocol (4). Matrix-assisted laser desorption ionization–time of flight (MALDI-TOF) mass spectrometry was used to identify the genus and species of the isolate, which was stored at -80°C until sequencing. From this freezer stock, *C. coyleae* UMB8490 was first streaked onto a Columbia nalidixic acid (CNA) agar plate and incubated at 35°C with 5% CO_2 for 24 h. A single colony was selected to inoculate LB broth and grown at 37°C with shaking for 24 h. DNA was extracted using the DNeasy blood and tissue kit following the manufacturer's protocol for Gram-positive bacteria with the following exceptions: we used $230\ \mu\text{l}$ of lysis buffer ($180\ \mu\text{l}$ 20 mM Tris-Cl, 2 mM sodium EDTA, and 1.2% Triton X-100 and $50\ \mu\text{l}$ lysozyme) in step 2 and altered the incubation time in step 5 to 10 min. DNA was quantified using the Qubit fluorometer and sent to the Microbial Genomic Sequencing Center (MiGS) at the University of Pittsburgh for sequencing, where the DNA was enzymatically fragmented using an Illumina tagmentation enzyme. Indices were attached using PCR and sequenced on the Illumina NextSeq 550 platform, producing 1,405,941 pairs of 150-bp reads. Raw reads were trimmed using Sickle v1.33 (<https://github.com/najoshi/sickle>) and assembled using SPAdes v3.13.0 with the “only-assembler” option for k values of 55, 77, 99, and 127 (7). Genome coverage was calculated using BBDMap v38.47 (<https://sourceforge.net/projects/bbmap/>). PATRIC v3.6.3 and RASTtk were used to annotate the genome sequences (8, 9), and the publicly available genome was annotated using the Prokaryotic Genome Annotation Pipeline (PGAP) v4.11 (10). Unless previously noted, default parameters were used for each software tool.

Citation Ali Khan A, Miller-Ensminger T, Voukadinova A, Wolfe AJ, Putonti C. 2020. Draft genome sequence of *Corynebacterium coyleae* UMB8490, isolated from the female urinary tract. Microbiol Resour Announc 9:e00393-20. <https://doi.org/10.1128/MRA.00393-20>.

Editor Frank J. Stewart, Georgia Institute of Technology

Copyright © 2020 Ali Khan et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Catherine Putonti, cputonti@luc.edu.

Received 12 April 2020

Accepted 19 April 2020

Published 14 May 2020

The *C. coyleae* UMB8490 draft genome is 2,483,223 bp long in 62 contigs with a GC content of 61.42%, genome coverage of 141.47×, and an N_{50} score of 92,141 bp. PGAP's annotation identified 2,290 protein-coding genes along with 51 tRNAs. PATRIC's annotation also identified a CRISPR array with 120 spacers. While PATRIC identified several genes associated with antibiotic resistance, analysis by ResFinder v3.2 (11) detected only resistance to aminoglycoside, which is used for single-dose treatment of UTIs (12). Further analysis of this genome will advance our understanding of the role that *C. coyleae* plays in the female urinary tract.

Data availability. This whole-genome shotgun project has been deposited in GenBank under the accession no. [JAAUVV000000000](https://www.ncbi.nlm.nih.gov/nuclseq/JAAUVV000000000). The version described in this paper is the first version, JAAUVV010000000. The raw sequencing reads have been deposited under the accession no. [SRR11441024](https://www.ncbi.nlm.nih.gov/nuclseq/SRR11441024).

ACKNOWLEDGMENTS

This work was conducted as part of Loyola University of Chicago's Department of Biology bacterial genomics course. For prior patient recruitment, we acknowledge the Loyola Urinary Education and Research Collaborative (LUEREC) and the patients who provided the samples for this study.

REFERENCES

- Bernard KA, Funke G. 2015. *Corynebacterium*, p 1–70. In Whitman WB, Rainey F, Kämpfer P, Trujillo M, Chun J, DeVos P, Hedlund B, Dedysh S (ed), *Bergey's manual of systematics of archaea and bacteria*. John Wiley & Sons, Ltd, Chichester, United Kingdom.
- Sokol-Leszczynska B, Leszczynski P, Lachowicz D, Rostkowska O, Niemczyk M, Piecha T, van Belkum A, Sawicka-Grzelak A, Mlynarczyk G. 2019. *Corynebacterium coyleae* as potential urinary tract pathogen. *Eur J Clin Microbiol Infect Dis* 38:1339–1342. <https://doi.org/10.1007/s10096-019-03565-4>.
- Pearce MM, Hilt EE, Rosenfeld AB, Zilliox MJ, Thomas-White K, Fok C, Kliethermes S, Schreckenberger PC, Brubaker L, Gai X, Wolfe AJ. 2014. The female urinary microbiome: a comparison of women with and without urgency urinary incontinence. *mBio* 5:e01283-14. <https://doi.org/10.1128/mBio.01283-14>.
- Hilt EE, McKinley K, Pearce MM, Rosenfeld AB, Zilliox MJ, Mueller ER, Brubaker L, Gai X, Wolfe AJ, Schreckenberger PC. 2014. Urine is not sterile: use of enhanced urine culture techniques to detect resident bacterial flora in the adult female bladder. *J Clin Microbiol* 52:871–876. <https://doi.org/10.1128/JCM.02876-13>.
- Jacobs KM, Thomas-White KJ, Hilt EE, Wolfe AJ, Waters TP. 2017. Microorganisms identified in the maternal bladder: discovery of the maternal bladder microbiota. *AJP Rep* 7:e188–e196. <https://doi.org/10.1055/s-0037-1606860>.
- Barberis CM, Montalvo E, Imas S, Traglia G, Almuzara MN, Rodriguez CH, Famiglietti A, Mazzocchi O, Vay C. 2018. Total nephrectomy following *Corynebacterium coyleae* urinary tract infection. *JMM Case Rep* 5:e005149. <https://doi.org/10.1099/jmmcr.0.005149>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyskhin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Brettin T, Davis JJ, Disz T, Edwards RA, Gerdes S, Olsen GJ, Olson R, Overbeek R, Parrello B, Pusch GD, Shukla M, Thomason JA, Stevens R, Vonstein V, Wattam AR, Xia F. 2015. RASTtk: a modular and extensible implementation of the RAST algorithm for building custom annotation pipelines and annotating batches of genomes. *Sci Rep* 5:8365. <https://doi.org/10.1038/srep08365>.
- Wattam AR, Davis JJ, Assaf R, Boisvert S, Brettin T, Bun C, Conrad N, Dietrich EM, Disz T, Gabbard JL, Gerdes S, Henry CS, Kenyon RW, Machi D, Mao C, Nordberg EK, Olsen GJ, Murphy-Olson DE, Olson R, Overbeek R, Parrello B, Pusch GD, Shukla M, Vonstein V, Warren A, Xia F, Yoo H, Stevens RL. 2017. Improvements to PATRIC, the all-bacterial Bioinformatics Database and Analysis Resource Center. *Nucleic Acids Res* 45:D535–D542. <https://doi.org/10.1093/nar/gkw1017>.
- Tatusova T, DiCuccio M, Badretdin A, Chetvermin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
- Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, Aarestrup FM, Larsen MV. 2012. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother* 67:2640–2644. <https://doi.org/10.1093/jac/dks261>.
- Goodlet KJ, Benhalima FZ, Nailor MD. 2018. A systematic review of single-dose aminoglycoside therapy for urinary tract infection: is it time to resurrect an old strategy? *Antimicrob Agents Chemother* 63:e02165–18. <https://doi.org/10.1128/AAC.02165-18>.