



5-14-2020

## Draft Genome Sequence of *Klebsiella pneumoniae* UMB8492, Isolated from the Female Urinary Tract

Suma Gondi  
*Loyola University Chicago*

Taylor Miller-Ensminger  
*Loyola University Chicago*

Adelina Voukadinova  
*Loyola University Chicago*

Alan J. Wolfe  
*Loyola University Chicago*, [awolfe@luc.edu](mailto:awolfe@luc.edu)

Catherine Putoni  
*Loyola University Chicago*, [cputonti@luc.edu](mailto:cputonti@luc.edu)

Follow this and additional works at: [https://ecommons.luc.edu/bioinformatics\\_facpub](https://ecommons.luc.edu/bioinformatics_facpub)



Part of the [Bioinformatics Commons](#), and the [Biology Commons](#)

### Recommended Citation

Gondi, Suma; Miller-Ensminger, Taylor; Voukadinova, Adelina; Wolfe, Alan J.; and Putoni, Catherine. Draft Genome Sequence of *Klebsiella pneumoniae* UMB8492, Isolated from the Female Urinary Tract. *Microbiology Resource Announcements*, 9, : , 2020. Retrieved from Loyola eCommons, Bioinformatics Faculty Publications, <http://dx.doi.org/10.1128/MRA.00405-2>

This Article is brought to you for free and open access by the Faculty Publications and Other Works by Department at Loyola eCommons. It has been accepted for inclusion in Bioinformatics Faculty Publications by an authorized administrator of Loyola eCommons. For more information, please contact [ecommons@luc.edu](mailto:ecommons@luc.edu).



This work is licensed under a [Creative Commons Attribution 4.0 License](#).  
© Gondi et al., 2020.



# Draft Genome Sequence of *Klebsiella pneumoniae* UMB8492, Isolated from the Female Urinary Tract

Suma Gondi,<sup>a</sup> Taylor Miller-Ensminger,<sup>b</sup> Adelina Voukadinova,<sup>b</sup> Alan J. Wolfe,<sup>c</sup>  Catherine Putonti<sup>a,b,c,d</sup>

<sup>a</sup>Department of Biology, Loyola University Chicago, Chicago, Illinois, USA

<sup>b</sup>Bioinformatics Program, Loyola University Chicago, Chicago, Illinois, USA

<sup>c</sup>Department of Microbiology and Immunology, Stritch School of Medicine, Loyola University Chicago, Maywood, Illinois, USA

<sup>d</sup>Department of Computer Science, Loyola University Chicago, Chicago, Illinois, USA

**ABSTRACT** Here, we present the draft genome sequence for *Klebsiella pneumoniae* strain UMB8492, isolated from a urine sample from a female patient with overactive bladder (OAB) symptoms. The assembled genome has a length of 5,660,575 bp, a G+C content of 57.19%, and an  $N_{50}$  value of 136,315 bp.

*Klebsiella pneumoniae* is a Gram-negative bacterium that is commonly found in the gastrointestinal tract of healthy animals and humans (1, 2). It can also cause urinary tract, respiratory tract, and bloodstream infections (3). *K. pneumoniae* is a major cause of hospital-acquired infections, and surveys from several different countries estimate that *K. pneumoniae* accounts for 2 to 6% of hospital-acquired UTIs (3, 4). *K. pneumoniae* is also classified as a multidrug-resistant (MDR) pathogen, and thus there are limited treatment options (1). Here, we present the draft genome of *K. pneumoniae* UMB8492, isolated from a catheterized urine sample. While *K. pneumoniae* is frequently associated with UTIs, the woman from whom it was isolated did not have a UTI but rather had symptoms of overactive bladder (OAB).

*K. pneumoniae* UMB8492 was isolated from a previously conducted institutional review board (IRB)-approved study (Loyola University Chicago, IRB no. LU207102) using the expanded quantitative urinary culture (EQUC) protocol (5). The genus and species of *Klebsiella pneumoniae* were confirmed using matrix-assisted laser desorption ionization–time of flight (MALDI-TOF) mass spectrometry before the culture was frozen at  $-80^{\circ}\text{C}$ . The EQUC method was also used to isolate strains of *Lactobacillus jensenii*, *Corynebacterium coyleae*, *Lactobacillus iners*, *Corynebacterium aurimucosum*, and *Staphylococcus epidermidis* from this same woman's urinary microbiome. From the freezer stock of UMB8492, an aliquot was streaked onto a Columbia nalidixic acid (CNA) agar plate and incubated at  $35^{\circ}\text{C}$  with 5%  $\text{CO}_2$  for 24 h. A single colony was isolated, and liquid LB medium was inoculated and grown overnight with shaking at  $37^{\circ}\text{C}$ . DNA was isolated from this liquid culture using the Qiagen DNeasy blood and tissue kit following the Gram-positive protocol with the following exceptions. We used 230  $\mu\text{l}$  of lysis buffer (180  $\mu\text{l}$  of 20 mM Tris-Cl, 2 mM sodium EDTA, and 1.2% Triton X-100 and 50  $\mu\text{l}$  of lysozyme) in step 2 and incubated the sample at  $56^{\circ}\text{C}$  for 10 minutes in step 5. The DNA quality was determined using a Qubit fluorometer. DNA was sent to the Microbial Genome Sequencing Center (MiGS) at the University of Pittsburgh for sequencing. MiGS prepared the libraries using the Illumina Nextera kit and sequenced the library on the NextSeq 550 platform. In total, 1,712,978 pairs of  $2 \times 150$ -bp reads were generated. These raw reads were trimmed using Sickle v1.33 (<https://github.com/najoshi/sickle>), using default parameters, and assembled using SPAdes v3.13.0 with the “only-assembler” option for k values of 55, 77, 99, and 127 (6). The genome assembly was annotated using the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) v4.11 (7).

**Citation** Gondi S, Miller-Ensminger T, Voukadinova A, Wolfe AJ, Putonti C. 2020. Draft genome sequence of *Klebsiella pneumoniae* UMB8492, isolated from the female urinary tract. Microbiol Resour Announc 9:e00405-20. <https://doi.org/10.1128/MRA.00405-20>.

**Editor** David A. Baltrus, University of Arizona

**Copyright** © 2020 Gondi et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Catherine Putonti, [cputonti@luc.edu](mailto:cputonti@luc.edu).

**Received** 13 April 2020

**Accepted** 27 April 2020

**Published** 14 May 2020

The genome coverage was calculated using BBMap v38.47 (<https://sourceforge.net/projects/bbmap/>) with default parameters.

*K. pneumoniae* UMB8492 has a draft genome size of 5,660,575 bp, assembled into 127 contigs with an  $N_{50}$  value of 136,315 bp, and a G+C content of 57.19%. The genome coverage is 77×. PGAP annotation identified 5,508 protein-coding genes and 79 tRNAs. This genome sequence provides an additional genome resource for the study of uropathogens.

**Data availability.** This whole-genome shotgun project has been deposited in GenBank under accession no. [JAAUVU000000000](https://ncbi.nlm.nih.gov/submit/submit.cgi?accession=JAAUVU000000000). The version described in this paper is the first version, JAAUVU010000000. The raw sequencing reads have been deposited in the SRA under accession no. [SRR11441023](https://www.ncbi.nlm.nih.gov/sra/SRR11441023).

## ACKNOWLEDGMENTS

This work was conducted as part of the Bacterial Genomics course at Loyola University Chicago's Department of Biology.

For prior patient recruitment, we acknowledge the Loyola Urinary Education and Research Collaborative (LUEREC) and the patients who provided the samples for this study.

## REFERENCES

1. Navon-Venezia S, Kondratyeva K, Carattoli A. 2017. *Klebsiella pneumoniae*: a major worldwide source and shuttle for antibiotic resistance. *FEMS Microbiol Rev* 41:252–275. <https://doi.org/10.1093/femsre/fux013>.
2. Wyres KL, Holt KE. 2018. *Klebsiella pneumoniae* as a key trafficker of drug resistance genes from environmental to clinically important bacteria. *Curr Opin Microbiol* 45:131–139. <https://doi.org/10.1016/j.mib.2018.04.004>.
3. Paczosa MK, Meccas J. 2016. *Klebsiella pneumoniae*: going on the offense with a strong defense. *Microbiol Mol Biol Rev* 80:629–661. <https://doi.org/10.1128/MMBR.00078-15>.
4. Meatherall BL, Gregson D, Ross T, Pitout JD, Laupland KB. 2009. Incidence, risk factors, and outcomes of *Klebsiella pneumoniae* bacteremia. *Am J Med* 122:866–873. <https://doi.org/10.1016/j.amjmed.2009.03.034>.
5. Hilt EE, McKinley K, Pearce MM, Rosenfeld AB, Zilliox MJ, Mueller ER, Brubaker L, Gai X, Wolfe AJ, Schreckenberger PC. 2014. Urine is not sterile: use of enhanced urine culture techniques to detect resident bacterial flora in the adult female bladder. *J Clin Microbiol* 52:871–876. <https://doi.org/10.1128/JCM.02876-13>.
6. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
7. Tatusova T, DiCuccio M, Badretdin A, Chetverin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.