



5-14-2020

## Draft Genome Sequence of *Lactobacillus jensenii* Strain UMB7766, Isolated from the Female Bladder

Taylor Miller-Ensminger  
*Loyola University Chicago*

Alan J. Wolfe  
*Loyola University Chicago*, [awolfe@luc.edu](mailto:awolfe@luc.edu)

Catherine Putoni  
*Loyola University Chicago*, [cputonti@luc.edu](mailto:cputonti@luc.edu)

Follow this and additional works at: [https://ecommons.luc.edu/bioinformatics\\_facpub](https://ecommons.luc.edu/bioinformatics_facpub)



Part of the [Bioinformatics Commons](#), and the [Biology Commons](#)

### Recommended Citation

Miller-Ensminger, Taylor; Wolfe, Alan J.; and Putoni, Catherine. Draft Genome Sequence of *Lactobacillus jensenii* Strain UMB7766, Isolated from the Female Bladder. *Microbiology Resource Announcements*, 9, : , 2020. Retrieved from Loyola eCommons, Bioinformatics Faculty Publications, <http://dx.doi.org/10.1128/MRA.00392-20>

This Article is brought to you for free and open access by the Faculty Publications and Other Works by Department at Loyola eCommons. It has been accepted for inclusion in Bioinformatics Faculty Publications by an authorized administrator of Loyola eCommons. For more information, please contact [ecommons@luc.edu](mailto:ecommons@luc.edu).



This work is licensed under a [Creative Commons Attribution 4.0 License](#).  
© Miller-Ensminger et al., 2020.



# Draft Genome Sequence of *Lactobacillus jensenii* Strain UMB7766, Isolated from the Female Bladder

Taylor Miller-Ensminger,<sup>a</sup> Alan J. Wolfe,<sup>b</sup>  Catherine Putonti<sup>a,b,c,d</sup>

<sup>a</sup>Bioinformatics Program, Loyola University Chicago, Chicago, Illinois, USA

<sup>b</sup>Department of Microbiology and Immunology, Stritch School of Medicine, Loyola University Chicago, Maywood, Illinois, USA

<sup>c</sup>Department of Biology, Loyola University Chicago, Chicago, Illinois, USA

<sup>d</sup>Department of Computer Science, Loyola University Chicago, Chicago, Illinois, USA

**ABSTRACT** *Lactobacillus jensenii* is a beneficial and prominent community member within both the vaginal and female urinary microbiota. As more genomes for *L. jensenii* strains are made publicly available, we gain more knowledge about this beneficial community member. Here, we present the draft genome sequence of *L. jensenii* UMB7766, which was isolated from a urine specimen from a catheterized female patient with recurrent urinary tract infections.

*Lactobacillus jensenii* is a dominant member of both the vaginal and female urinary microbiota of healthy individuals (1, 2). *L. jensenii* acts as a protective species within both niches, showing bactericidal effects against pathogens. Notably, in the urinary tract, *L. jensenii* has been shown to reduce the growth of uropathogenic *Escherichia coli* (3), the most common cause of infrequent urinary tract infections (4–6). Last year, our group released 11 *L. jensenii* genomes (7), raising the number of characterized *L. jensenii* strains from the bladder to 13. To add to this collection, we present findings for *L. jensenii* UMB7766, which was isolated from a urine specimen from a catheterized female patient with recurrent urinary tract infections.

The sample was collected from a patient at the Women's Pelvic Medicine Center at the University of California, San Diego, California, in August 2017 as part of an institutional review board (IRB)-approved study (University of California, San Diego, IRB approval number 170077AW). Urine was cultured using the expanded quantitative urine culture (EQUC) method (8). The genus and species for *L. jensenii* UMB7766 were determined by matrix-assisted laser desorption ionization–time of flight (MALDI-TOF) mass spectrometry following the protocol detailed previously (8). The strain was streaked onto a Columbia nalidixic acid (CNA) agar plate with 5% sheep blood plate (catalog number 221353; BD). The plate was incubated at 35°C in 5% CO<sub>2</sub> for 48 h. A single colony was selected and grown in liquid MRS medium supplemented with 1 ml/liter Tween 80. DNA was extracted using a blood and tissue kit (catalog number 69504; Qiagen), following the manufacturer's protocol for Gram-positive organisms, and quantified using a Qubit fluorometer. Library preparation and sequencing were performed by the Microbial Genome Sequencing Center at the University of Pittsburgh. Library preparation was performed using Illumina Nextera chemistry, with sequencing on the Illumina NextSeq 550 platform (2 × 150-bp reads). Sequencing produced 3,314,892 pairs of reads. Raw reads were trimmed using Sickle v1.33 (<https://github.com/najoshi/sickle>) and assembled using SPAdes v3.13.0 with the only-assembler option for k values of 55, 77, 99, and 127 (9). The coverage was calculated using BBMap v38.47 (<https://sourceforge.net/projects/bbmap>). The genome was annotated using the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) v4.11 (10). The CRISPRCasFinder Web server was used to detect CRISPR

**Citation** Miller-Ensminger T, Wolfe AJ, Putonti C. 2020. Draft genome sequence of *Lactobacillus jensenii* strain UMB7766, isolated from the female bladder. Microbiol Resour Announc 9:e00392-20. <https://doi.org/10.1128/MRA.00392-20>.

**Editor** Irene L. G. Newton, Indiana University, Bloomington

**Copyright** © 2020 Miller-Ensminger et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Catherine Putonti, [cputonti@luc.edu](mailto:cputonti@luc.edu).

**Received** 12 April 2020

**Accepted** 24 April 2020

**Published** 14 May 2020

arrays (11), and Mash was used to identify similar genomes (12) through the PATRIC v3.6.3 Web server (13). Unless specified otherwise, default parameters were used for all software tools.

The *L. jensenii* UMB7766 genome is 1,710,714 bp long. It was assembled into 41 contigs with an  $N_{50}$  value of 68,917 bp and  $475\times$  coverage. At 34.32%, the GC content of this strain is similar to that of other *L. jensenii* strains available in GenBank. PGAP identified 1,637 genes, with 1,529 protein-coding sequences. Furthermore, it identified 54 tRNA sequences, 3 5S rRNA sequences, 1 16S rRNA sequence, and 1 23S rRNA sequence. A type II CRISPR system was identified, and this strain has 1 CRISPR array with 15 spacers. The complete genome of *L. jensenii* UMB7766 was also compared to those of all publicly available strains of this species, including other isolates from the urinary tract and isolates from the vaginal and gut microbiota. This comparison identified *L. jensenii* UMB3442 (whole-genome sequencing GenBank accession number [VYVY00000000](https://doi.org/10.1128/JB.01439-13)), which was also isolated from the female urinary microbiota, as having the closest related genome. This finding suggests that *L. jensenii* strains of the urinary tract may more closely resemble each other than those of the vaginal microbiota.

**Data availability.** The whole-genome sequencing project for *L. jensenii* UMB7766 has been deposited in GenBank under the accession number [JAAUWC00000000](https://doi.org/10.1128/JA.00000000). The raw sequencing reads have been deposited in GenBank under the accession number [SRR11441033](https://doi.org/10.1128/SRR11441033).

## ACKNOWLEDGMENTS

This work was conducted as part of the Loyola University Chicago's Department of Biology Bacterial Genomics course. For prior patient recruitment, we acknowledge the Loyola Urinary Education and Research Collaborative and the patients who provided the samples for this study.

## REFERENCES

- Mendes-Soares H, Suzuki H, Hickey RJ, Forney LJ. 2014. Comparative functional genomics of *Lactobacillus* spp. reveals possible mechanisms for specialization of vaginal lactobacilli to their environment. *J Bacteriol* 196:1458–1470. <https://doi.org/10.1128/JB.01439-13>.
- Thomas-White K, Forster SC, Kumar N, Van Kuiken M, Putonti C, Stares MD, Hilt EE, Price TK, Wolfe AJ, Lawley TD. 2018. Culturing of female bladder bacteria reveals an interconnected urogenital microbiota. *Nat Commun* 9:1557. <https://doi.org/10.1038/s41467-018-03968-5>.
- Kalyoussef S, Nieves E, Dinerman E, Carpenter C, Shankar V, Oh J, Burd B, Angeletti RH, Buckheit KW, Fredricks DN, Madan RP, Keller MJ, Herold BC. 2012. *Lactobacillus* proteins are associated with the bactericidal activity against *E. coli* of female genital tract secretions. *PLoS One* 7:e49506. <https://doi.org/10.1371/journal.pone.0049506>.
- Foxman B. 2014. Urinary tract infection syndromes: occurrence, recurrence, bacteriology, risk factors, and disease burden. *Infect Dis Clin North Am* 28:1–13. <https://doi.org/10.1016/j.idc.2013.09.003>.
- Flores-Mireles AL, Walker JN, Caparon M, Hultgren SJ. 2015. Urinary tract infections: epidemiology, mechanisms of infection and treatment options. *Nat Rev Microbiol* 13:269–284. <https://doi.org/10.1038/nrmicro3432>.
- Zhang L, Foxman B, Marrs C. 2002. Both urinary and rectal *Escherichia coli* isolates are dominated by strains of phylogenetic group B2. *J Clin Microbiol* 40:3951–3955. <https://doi.org/10.1128/jcm.40.11.3951-3955.2002>.
- Putonti C, Ahmad A, Baddoo G, Diaz J, Do M, Gallian N, Lorentzen C, Mohammed H, Murphy J, Olu-Ajeigbe A, Yang T, Miller-Ensminger T, Stark N, Maskeri L, Van Dusen J, Wolfe AJ. 2019. Draft genome sequences of 11 *Lactobacillus jensenii* strains isolated from the female bladder. *Microbiol Resour Announc* 8:e00970-19. <https://doi.org/10.1128/MRA.00970-19>.
- Hilt EE, McKinley K, Pearce MM, Rosenfeld AB, Zilliox MJ, Mueller ER, Brubaker L, Gai X, Wolfe AJ, Schreckenberger PC. 2014. Urine is not sterile: use of enhanced urine culture techniques to detect resident bacterial flora in the adult female bladder. *J Clin Microbiol* 52:871–876. <https://doi.org/10.1128/JCM.02876-13>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
- Grissa I, Vergnaud G, Pourcel C. 2007. CRISPRFinder: a Web tool to identify clustered regularly interspaced short palindromic repeats. *Nucleic Acids Res* 35:W52–W57. <https://doi.org/10.1093/nar/gkm360>.
- Ondov BD, Treangen TJ, Melsted P, Mallonee AB, Bergman NH, Koren S, Phillippy AM. 2016. Mash: fast genome and metagenome distance estimation using MinHash. *Genome Biol* 17:132. <https://doi.org/10.1186/s13059-016-0997-x>.
- Wattam AR, Davis JJ, Assaf R, Boisvert S, Brettin T, Bun C, Conrad N, Dietrich EM, Disz T, Gabbard JL, Gerdes S, Henry CS, Kenyon RW, Machi D, Mao C, Nordberg EK, Olsen GJ, Murphy-Olson DE, Olson R, Overbeek R, Parrello B, Pusch GD, Shukla M, Vonstein V, Warren A, Xia F, Yoo H, Stevens RL. 2017. Improvements to PATRIC, the all-bacterial Bioinformatics Database and Analysis Resource Center. *Nucleic Acids Res* 45:D535–D542. <https://doi.org/10.1093/nar/gkw1017>.