



5-14-2020

## Draft Genome Sequence of *Streptococcus anginosus* UMB1296, Isolated from the Female Urinary Tract

Sara Temelkova  
*Loyola University Chicago*

Taylor Miller-Ensminger  
*Loyola University Chicago*

Adelina Voukadinova  
*Loyola University Chicago*

Alan J. Wolfe  
*Loyola University Chicago*, [awolfe@luc.edu](mailto:awolfe@luc.edu)

Catherine Putoni  
*Loyola University Chicago*, [cputonti@luc.edu](mailto:cputonti@luc.edu)

Follow this and additional works at: [https://ecommons.luc.edu/bioinformatics\\_facpub](https://ecommons.luc.edu/bioinformatics_facpub)



Part of the [Bioinformatics Commons](#), and the [Biology Commons](#)

### Recommended Citation

Temelkova, Sara; Miller-Ensminger, Taylor; Voukadinova, Adelina; Wolfe, Alan J.; and Putoni, Catherine. Draft Genome Sequence of *Streptococcus anginosus* UMB1296, Isolated from the Female Urinary Tract. *Microbiology Resource Announcements*, 9, : , 2020. Retrieved from Loyola eCommons, Bioinformatics Faculty Publications, <http://dx.doi.org/10.1128/MRA.00409-20>

This Article is brought to you for free and open access by the Faculty Publications and Other Works by Department at Loyola eCommons. It has been accepted for inclusion in Bioinformatics Faculty Publications by an authorized administrator of Loyola eCommons. For more information, please contact [ecommons@luc.edu](mailto:ecommons@luc.edu).



This work is licensed under a [Creative Commons Attribution 4.0 License](#).  
© Temelkova et al., 2020.



# Draft Genome Sequence of *Streptococcus anginosus* UMB1296, Isolated from the Female Urinary Tract

Sara Temelkova,<sup>a</sup> Taylor Miller-Ensminger,<sup>b</sup> Adelina Voukadinova,<sup>b</sup> Alan J. Wolfe,<sup>c</sup>  Catherine Putonti<sup>b,c,d,e</sup>

<sup>a</sup>Neuroscience Program, Loyola University Chicago, Chicago, Illinois, USA

<sup>b</sup>Bioinformatics Program, Loyola University Chicago, Chicago, Illinois, USA

<sup>c</sup>Department of Microbiology and Immunology, Stritch School of Medicine, Loyola University Chicago, Maywood, Illinois, USA

<sup>d</sup>Department of Biology, Loyola University Chicago, Chicago, Illinois, USA

<sup>e</sup>Department of Computer Science, Loyola University Chicago, Chicago, Illinois, USA

**ABSTRACT** We present the draft genome sequence of a *Streptococcus anginosus* strain isolated from the female urinary tract. The *S. anginosus* UMB1296 draft genome has a size of 1,924,009 bp assembled into 35 contigs with a GC content of 38.69%. Genome annotation revealed 1,775 protein-coding genes, including several known virulence factors.

Though universal consensus on its taxonomy is yet to be established, *Streptococcus anginosus* is commonly classified as part of the *Streptococcus milleri* group (SMG) of the genus *Streptococcus* (1, 2). Members of the SMG, which includes *S. anginosus*, *S. intermedius*, and *S. constellatus*, belong to the natural flora of human mucous membranes and healthy female urogenital tracts (2). However, they are known for their association with purulent infections throughout the body and distinct ability for causing abscesses (2, 3). Of the SMG species, *S. anginosus* has been most frequently identified from genitourinary sources (4, 5) and is largely shaped by virulence traits (6). Investigation of infections caused by *S. anginosus* have been limited, and as a result, its pathogenic potential has been historically underrecognized (7).

*S. anginosus* UMB1296 was obtained from a catheterized urine sample from a female with a urinary tract infection. The sample was clinically isolated using the expanded quantitative urinary (EQUC) protocol (8) from a prior institutional review board (IRB)-approved study (9). The genus and species for this isolate were determined via matrix-assisted laser desorption ionization–time of flight (MALDI-TOF) mass spectrometry following a previously described protocol (8). The isolate was then stored at  $-80^{\circ}\text{C}$  until sequencing. The *S. anginosus* sample was streaked onto a Columbia nalidixic acid agar plate using the quadrant streaking method and a sterile inoculating loop. The plate was incubated for 24 h at  $35^{\circ}\text{C}$  with 5%  $\text{CO}_2$ . A sterile inoculating loop was used to isolate one colony into 1 ml *Actinomyces* broth (catalog no. 40834; Millipore), and the culture was grown overnight under the same conditions as before. DNA was extracted using the Qiagen DNeasy blood and tissue kit with a modified Gram-positive extraction protocol to include the addition of 230  $\mu\text{l}$  of lysis buffer (180  $\mu\text{l}$  of 20 mM Tris-Cl, 2 mM sodium EDTA, and 1.2% Triton X-100 and 50  $\mu\text{l}$  of lysozyme) to the culture pellet and incubation at  $56^{\circ}\text{C}$  for 10 min in a mixture of 25  $\mu\text{l}$  of proteinase K and 200  $\mu\text{l}$  of buffer AL. The extracted DNA was quantified using a Qubit fluorometer. DNA was sent to the Microbial Genomic Sequencing Center (MiGS) at the University of Pittsburgh for sequencing, where the DNA was first enzymatically fragmented into indices using an Illumina tagmentation enzyme. Indices were attached using PCR and sequenced using an Illumina NextSeq 550 flow cell, producing 1,765,717 pairs of paired-end reads 150 bp long. The raw reads were trimmed using Sickle v1.33 (<https://github.com/najoshi/sickle>)

**Citation** Temelkova S, Miller-Ensminger T, Voukadinova A, Wolfe AJ, Putonti C. 2020. Draft genome sequence of *Streptococcus anginosus* UMB1296, isolated from the female urinary tract. *Microbiol Resour Announc* 9:e00409-20. <https://doi.org/10.1128/MRA.00409-20>.

**Editor** Simon Roux, DOE Joint Genome Institute

**Copyright** © 2020 Temelkova et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Catherine Putonti, [cputonti@luc.edu](mailto:cputonti@luc.edu).

**Received** 13 April 2020

**Accepted** 20 April 2020

**Published** 14 May 2020

and assembled using SPAdes v3.13.0 with the “only-assembler” option for k values of 55, 77, 99, and 127 (10). Genome coverage was calculated using BBMap v38.47 (<https://sourceforge.net/projects/bbmap/>). While PATRIC v3.6.6 (11) was used initially to annotate the genome sequences, the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) v4.11 (12) was used for reannotation and accompanies the publicly available genome. Unless previously noted, default parameters were used for each software tool.

The *S. anginosus* UMB1296 draft genome assembly is 1,924,009 bp long and is assembled into 35 contigs, with a genome coverage of 235×, a GC content of 38.69%, and an  $N_{50}$  score of 122,329 bp. The PGAP annotation includes 1,775 protein-coding genes. PATRIC identified 59 virulence factors, including those previously identified for the species (6), e.g., the pneumococcal surface adhesion protein PsaA. Due to the known role of the urinary microbiota in a variety of genitourinary-associated disorders, characterizing the members of this microbial community allows for further understanding of associated clinical conditions.

**Data availability.** This whole-genome shotgun project has been deposited in GenBank under the accession no. [JAAUWH00000000](https://www.ncbi.nlm.nih.gov/GenBank/JAAUWH00000000). The version described in this paper is the first version, JAAUWH010000000. The raw sequencing reads have been deposited in SRA under the accession no. [SRR11441015](https://www.ncbi.nlm.nih.gov/SRA/SRR11441015).

## ACKNOWLEDGMENTS

This work was conducted as part of the Bacterial Genomics course at Loyola University Chicago’s Department of Biology. For prior patient recruitment, we acknowledge the Loyola Urinary Education and Research Collaborative (LUEREC) and the patients who provided the samples for this study.

## REFERENCES

- Jensen A, Hoshino T, Kilian M. 2013. Taxonomy of the Anginosus group of the genus *Streptococcus* and description of *Streptococcus anginosus* subsp. *whileyi* subsp. nov. and *Streptococcus constellatus* subsp. *viborgensis* subsp. nov. *Int J Syst Evol Microbiol* 63:2506–2519. <https://doi.org/10.1099/ijs.0.043232-0>.
- Grinwis ME, Sibley CD, Parkins MD, Eshaghrushan CS, Rabin HR, Surette MG. 2010. Characterization of *Streptococcus milleri* group isolates from expectorated sputum of adult patients with cystic fibrosis. *J Clin Microbiol* 48:395–401. <https://doi.org/10.1128/JCM.01807-09>.
- Doern CD, Burnham C-A. 2010. It’s not easy being green: the viridans group streptococci, with a focus on pediatric clinical manifestations. *J Clin Microbiol* 48:3829–3835. <https://doi.org/10.1128/JCM.01563-10>.
- Whiley RA, Beighton D, Winstanley TG, Fraser HY, Hardie JM. 1992. *Streptococcus intermedius*, *Streptococcus constellatus*, and *Streptococcus anginosus* (the *Streptococcus milleri* group): association with different body sites and clinical infections. *J Clin Microbiol* 30:243–244. <https://doi.org/10.1128/JCM.30.1.243-244.1992>.
- Furuichi M, Horikoshi Y. 2018. Sites of infection associated with *Streptococcus anginosus* group among children. *J Infect Chemother* 24:99–102. <https://doi.org/10.1016/j.jiac.2017.09.011>.
- Sitkiewicz I. 2018. How to become a killer, or is it all accidental? Virulence strategies in oral streptococci. *Mol Oral Microbiol* 33:1–12. <https://doi.org/10.1111/omi.12192>.
- Asam D, Spellerberg B. 2014. Molecular pathogenicity of *Streptococcus anginosus*. *Mol Oral Microbiol* 29:145–155. <https://doi.org/10.1111/omi.12056>.
- Hilt EE, McKinley K, Pearce MM, Rosenfeld AB, Zilliox MJ, Mueller ER, Brubaker L, Gai X, Wolfe AJ, Schreckenberger PC. 2014. Urine is not sterile: use of enhanced urine culture techniques to detect resident bacterial flora in the adult female bladder. *J Clin Microbiol* 52:871–876. <https://doi.org/10.1128/JCM.02876-13>.
- Price TK, Dune T, Hilt EE, Thomas-White KJ, Kliethermes S, Brincat C, Brubaker L, Wolfe AJ, Mueller ER, Schreckenberger PC. 2016. The clinical urine culture: enhanced techniques improve detection of clinically relevant microorganisms. *J Clin Microbiol* 54:1216–1222. <https://doi.org/10.1128/JCM.00044-16>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Brettin T, Davis JJ, Disz T, Edwards RA, Gerdes S, Olsen GJ, Olson R, Overbeek R, Parrello B, Pusch GD, Shukla M, Thomason JA, III, Stevens R, Vonstein V, Wattam AR, Xia F. 2015. RASTtk: a modular and extensible implementation of the RAST algorithm for building custom annotation pipelines and annotating batches of genomes. *Sci Rep* 5:8365. <https://doi.org/10.1038/srep08365>.
- Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.