



1-2-2020

Complete Genome Sequences of Streptococcus mitis Strains Isolated from the Oral Cavity and Urogenital Tract of a Woman and Her Male Sexual Partner

Carine R. Mores
Loyola University Chicago

Travis K. Price
Loyola University Chicago

Bridget Brassil
Loyola University Chicago

Catherine Putoni
Loyola University Chicago, cputonti@luc.edu

Alan J. Wolfe
Loyola University Chicago, awolfe@luc.edu

Follow this and additional works at: https://ecommons.luc.edu/bioinformatics_facpub

 Part of the [Bioinformatics Commons](#), and the [Biology Commons](#)

Recommended Citation

Mores, Carine R.; Price, Travis K.; Brassil, Bridget; Putoni, Catherine; and Wolfe, Alan J.. Complete Genome Sequences of Streptococcus mitis Strains Isolated from the Oral Cavity and Urogenital Tract of a Woman and Her Male Sexual Partner. *Microbiology Resource Announcements*, 9, : , 2020. Retrieved from Loyola eCommons, Bioinformatics Faculty Publications, <http://dx.doi.org/10.1128/MRA.01379-19>

This Article is brought to you for free and open access by the Faculty Publications and Other Works by Department at Loyola eCommons. It has been accepted for inclusion in Bioinformatics Faculty Publications by an authorized administrator of Loyola eCommons. For more information, please contact ecommons@luc.edu.



This work is licensed under a [Creative Commons Attribution 4.0 License](#).
© Mores et al., 2020.



Complete Genome Sequences of *Streptococcus mitis* Strains Isolated from the Oral Cavity and Urogenital Tract of a Woman and Her Male Sexual Partner

Carine R. Mores,^{a,b} Travis K. Price,^a Bridget Brassil,^b  Catherine Putonti,^{a,b,c,d} Alan J. Wolfe^a

^aDepartment of Microbiology and Immunology, Stritch School of Medicine, Loyola University Chicago, Maywood, Illinois, USA

^bDepartment of Biology, Loyola University Chicago, Chicago, Illinois, USA

^cBioinformatics Program, Loyola University Chicago, Chicago, Illinois, USA

^dDepartment of Computer Science, Loyola University Chicago, Chicago, Illinois, USA

ABSTRACT *Streptococcus mitis* is a member of the mitis group of the genus *Streptococcus*, which includes commensal species of the oral cavity and upper respiratory tract. Here, we report 39 complete genome sequences of *S. mitis* strains isolated from the oral cavity and urogenital tract of a woman and her male sexual partner.

Streptococcus mitis is an abundant member of the commensal microbiota of the upper respiratory tract and oral cavity (1, 2). Although it is considered to be a commensal species, *S. mitis* can cause a variety of invasive diseases in human (3). In order to increase our knowledge of this commensal bacterium, we isolated multiple strains of *S. mitis* from a woman and her male sexual partner. These strains were isolated from different anatomical sites and on different days. Here, we present the genome sequences for a subset of this collection.

Samples were collected from oral swabs, vaginal swabs, periurethral swabs, penile swabs, and voided urine samples from one female and her male sexual partner as part of an institutional review board (IRB)-approved study (LU 209830). Swabs were collected using the BD liquid Amies elution swab (ESwab) collection system. Urine samples were collected as clean-catch midstream voided urine. (Note that the participant was given instructions for obtaining the voided urine samples, and the periurethral swab and urine samples differed significantly [4].) Strains were isolated using a modified version of the expanded quantitative urine culture (EQUC) protocol (5) and stored at -80°C . From these samples, 39 *S. mitis* strains, identified by matrix-assisted laser desorption ionization–time of flight (MALDI-TOF) mass spectrometry, were selected for whole-genome sequencing. Each *S. mitis* isolate was grown on Columbia colistin-naladixic acid agar with 5% sheep blood plates incubated under 5% CO_2 conditions at 37°C for 48 h. Pure cultures of *S. mitis* were transferred to brain heart infusion (BHI) broth with 10% fetal bovine serum (FBS) and incubated under 5% CO_2 conditions at 37°C for 48 h and pelleted. DNA was extracted from pellets using a phenol-chloroform method and quantified using a Qubit 2.0 fluorometer and an Agilent Bioanalyzer. DNA libraries were created using the Illumina Nextera kit and sequenced using the MiSeq reagent kit v2, producing, on average, 336,117 pairs of 250-bp reads. Quality control and demultiplexing of sequence data were done with onboard MiSeq Control software and MiSeq Reporter v3.1. Raw reads were trimmed using Sickle v1.33 (<https://github.com/najoshi/sickle>) and assembled using SPAdes v3.13.0 (6) with the “only-assembler” option for k values of 55, 77, 99, and 127. Genome coverage was calculated using BMap v38.47 (<https://sourceforge.net/projects/bbmap/>). The NCBI Prokaryotic Genome Annotation Pipeline (PGAP) v4.8 (7) was used to annotate the genome sequences. Unless previously noted, default parameters were used for each software tool.

Citation Mores CR, Price TK, Brassil B, Putonti C, Wolfe AJ. 2020. Complete genome sequences of *Streptococcus mitis* strains isolated from the oral cavity and urogenital tract of a woman and her male sexual partner. Microbiol Resour Announc 9:e01379-19. <https://doi.org/10.1128/MRA.01379-19>.

Editor Julie C. Dunning Hotopp, University of Maryland School of Medicine

Copyright © 2020 Mores et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Catherine Putonti, cputonti@luc.edu.

Received 3 November 2019

Accepted 19 November 2019

Published 2 January 2020

TABLE 1 Genome assembly and annotation statistics

Strain	Partner sex	Isolation source	Coverage (×)	No. of contigs	N_{50} (bp)	Genome length (bp)	No. of CDS	No. of rRNAs	No. of tRNAs	GenBank WGS accession no. ^a	SRA accession no.
SM05	Female	Voided urine sample	59.22	23	194,401	2,001,354	1,950	3	44	WIKE000000000	SRR10341581
SM07	Female	Periurethral swab	64.74	27	199,509	2,002,948	1,956	3	44	WIKD000000000	SRR10341580
SM09	Female	Oral swab	68.4	15	187,601	1,973,655	1,962	3	43	WIKC000000000	SRR10341605
SM12	Male	Oral swab	33.68	46	105,495	2,095,125	2,043	3	42	WIKB000000000	SRR10341594
SM20	Female	Vaginal swab	50.33	55	82,209	2,008,422	1,969	3	43	WIKA000000000	SRR10341584
SM26	Female	Oral swab	58.51	27	251,708	2,060,956	1,982	3	41	WIJZ000000000	SRR10341583
SM27	Female	Voided urine sample	53.29	27	174,586	2,001,505	1,957	3	44	WIJY000000000	SRR10341582
SM30	Female	Periurethral swab	51.27	28	199,508	2,007,955	1,965	3	44	WIJX000000000	SRR10341577
SM33	Male	Oral swab	15.19	55	74,863	2,093,429	2,047	3	39	WIJW000000000	SRR10341576
SM39	Female	Oral swab	46.12	127	27,218	2,015,490	2,042	3	47	WIJV000000000	SRR10341575
SM45	Female	Voided urine sample	44.13	25	148,327	2,002,001	1,953	3	44	WIJU000000000	SRR10341579
SM48	Male	Oral swab	19.61	45	105,607	2,094,194	2,038	3	42	WIJT000000000	SRR10341578
SM04	Female	Voided urine sample	51.52	21	225,775	2,000,773	1,956	3	44	WIJS000000000	SRR10341613
SM06	Female	Voided urine sample	59.03	27	188,320	2,002,707	1,957	3	44	WIJR000000000	SRR10341612
SM08	Female	Periurethral swab	51.75	18	205,523	1,996,012	1,960	3	44	WIJQ000000000	SRR10341611
SM10	Female	Oral swab	51.35	49	132,942	2,168,718	2,121	4	39	WIJP000000000	SRR10341610
SM11	Female	Oral swab	56.8	113	30,148	2,014,391	2,031	4	47	WIJO000000000	SRR10341609
SM13	Male	Oral swab	48.15	120	97,578	2,152,221	2,150	3	43	WIJN000000000	SRR10341608
SM14	Male	Oral swab	47.22	58	64,862	2,094,407	2,055	3	42	WIJM000000000	SRR10341607
SM15	Male	Penile swab	44.11	21	155,584	2,016,502	2,018	3	52	WIJL000000000	SRR10341606
SM16	Male	Penile swab	53.16	67	72,040	2,017,910	1,886	2	40	WIJK000000000	SRR10341604
SM28	Female	Voided urine sample	59.9	30	186,870	2,007,541	1,963	3	44	WIJJ000000000	SRR10341603
SM29	Female	Voided urine sample	47.58	24	199,508	2,005,875	1,961	3	44	WIJI000000000	SRR10341602
SM34	Male	Oral swab	18.72	46	105,662	2,091,923	2,047	3	43	WIJH000000000	SRR10341601
SM40	Female	Oral swab	53.86	124	27,218	2,015,048	2,039	3	47	WIJG000000000	SRR10341600
SM41	Female	Oral swab	46.7	27	177,356	2,032,459	2,058	3	44	WIJF000000000	SRR10341599
SM46	Female	Voided urine sample	52.34	28	187,795	2,027,642	1,964	3	44	WIJE000000000	SRR10341598
SM47	Female	Voided urine sample	46.19	31	147,586	2,006,587	1,965	3	44	WIJD000000000	SRR10341597
SM49	Male	Oral swab	64.52	30	141,856	1,996,076	1,958	2	44	WIJC000000000	SRR10341596
SM50	Male	Oral swab	49.28	18	208,384	1,894,166	1,859	4	49	WIJB000000000	SRR10341595
SM17	Female	Voided urine sample	33.99	450	7,154	1,984,878	1,944	2	37	WIJA000000000	SRR10341593
SM18	Female	Voided urine sample	137.48	30	147,586	2,000,922	1,954	3	44	WIIZ000000000	SRR10341592
SM19	Female	Voided urine sample	111.07	27	186,870	2,006,191	1,965	3	44	WIY000000000	SRR10341591
SM35	Female	Voided urine sample	100.62	25	187,601	2,007,565	1,963	3	44	WIIX000000000	SRR10341590
SM36	Female	Voided urine sample	106	25	186,870	2,005,820	1,964	3	44	WIIW000000000	SRR10341589
SM37	Female	Voided urine sample	76.02	29	188,320	2,006,041	1,963	3	44	WIIV000000000	SRR10341588
SM42	Female	Voided urine sample	115.16	33	131,530	2,007,726	1,962	3	44	WIIU000000000	SRR10341587
SM43	Female	Voided urine sample	94.02	26	147,586	2,000,668	1,954	3	44	WIIT000000000	SRR10341586
SM44	Female	Voided urine sample	115.16	23	186,871	2,000,125	1,953	3	44	WIIS000000000	SRR10341585

^aWGS, whole-genome shotgun.

Table 1 lists all of the 39 *S. mitis* strains and their source, as well as their genome assembly statistics. The average GC content is 41%, similar to that reported in GenBank for other strains of the species. Annotations identified an average of 1,985 coding sequences (CDS) (Table 1). The strains varied in their numbers of rRNA operons and tRNAs. The addition here of 39 genome sequences greatly increases our knowledge of the genetic diversity of this bacterial species within the oral and lower urinary tract microbiota.

Data availability. This whole-genome shotgun (WGS) project has been deposited in GenBank, and the accession numbers for each genome assembly are listed in Table 1. The versions described in this paper are the first versions. Raw sequence data are publicly available in SRA for the 39 *S. mitis* strains; the accession numbers are listed in Table 1. The WGS and SRA records are associated with BioProject number [PRJNA316969](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA316969).

ACKNOWLEDGMENTS

We acknowledge The Loyola Urinary Education and Research Collaborative for recruitment of study participants and attainment of urine samples and Roberto Limeira of the Loyola Genomics Facility for sequencing of the bacterial genomes.

REFERENCES

1. Aas JA, Paster BJ, Stokes LN, Olsen I, Dewhirst FE. 2005. Defining the normal bacterial flora of the oral cavity. *J Clin Microbiol* 43:5721–5732. <https://doi.org/10.1128/JCM.43.11.5721-5732.2005>.
2. Frandsen EV, Pedrazzoli V, Kilian M. 1991. Ecology of viridans streptococci in the oral cavity and pharynx. *Oral Microbiol Immunol* 6:129–133. <https://doi.org/10.1111/j.1399-302x.1991.tb00466.x>.
3. Mitchell J. 2011. *Streptococcus mitis*: walking the line between commensalism and pathogenesis. *Mol Oral Microbiol* 26:89–98. <https://doi.org/10.1111/j.2041-1014.2010.00601.x>.
4. Wolff BJ, Price TK, Joyce CJ, Wolfe AJ, Mueller ER. 2019. Oral probiotics and the female urinary microbiome: a double-blinded randomized placebo-controlled trial. *Int Urol Nephrol* 51:2149–2159. <https://doi.org/10.1007/s11255-019-02282-3>.
5. Price TK, Dune T, Hilt EE, Thomas-White KJ, Kliethermes S, Brincat C, Brubaker L, Wolfe AJ, Mueller ER, Schreckenberger PC. 2016. The clinical urine culture: enhanced techniques improve detection of clinically relevant microorganisms. *J Clin Microbiol* 54:1216–1222. <https://doi.org/10.1128/JCM.00044-16>.
6. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
7. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44: 6614–6624. <https://doi.org/10.1093/nar/gkw569>.