



6-4-2020

Draft Genome Sequence of *Staphylococcus epidermidis* UMB8493, Isolated from the Female Urinary Tract

Anthony Truckenbrod
Loyola University Chicago

Taylor Miller-Ensminger
Loyola University Chicago

Adelina Voukadinova
Loyola University Chicago

Alan J. Wolfe
Loyola University Chicago, awolfe@luc.edu

Catherine Putonti
Loyola University Chicago, cputonti@luc.edu

Follow this and additional works at: https://ecommons.luc.edu/bioinformatics_facpub

 Part of the [Bioinformatics Commons](#)

Recommended Citation

Truckenbrod, Anthony; Miller-Ensminger, Taylor; Voukadinova, Adelina; Wolfe, Alan J.; and Putonti, Catherine. Draft Genome Sequence of *Staphylococcus epidermidis* UMB8493, Isolated from the Female Urinary Tract. *Microbiology Resource Announcements*, 9, : , 2020. Retrieved from Loyola eCommons, Bioinformatics Faculty Publications, <http://dx.doi.org/10.1128/MRA.00419-20>

This Article is brought to you for free and open access by the Faculty Publications and Other Works by Department at Loyola eCommons. It has been accepted for inclusion in Bioinformatics Faculty Publications by an authorized administrator of Loyola eCommons. For more information, please contact ecommons@luc.edu.



This work is licensed under a [Creative Commons Attribution 4.0 License](#).
© Truckenbrod et al., 2020.



Draft Genome Sequence of *Staphylococcus epidermidis* UMB8493, Isolated from the Female Urinary Tract

Anthony Truckenbrod,^a Taylor Miller-Ensminger,^b Adelina Voukadinova,^b Alan J. Wolfe,^c  Catherine Putonti^{a,b,c,d}

^aDepartment of Biology, Loyola University Chicago, Chicago, Illinois, USA

^bBioinformatics Program, Loyola University Chicago, Chicago, Illinois, USA

^cDepartment of Microbiology and Immunology, Stritch School of Medicine, Loyola University Chicago, Maywood, Illinois, USA

^dDepartment of Computer Science, Loyola University Chicago, Chicago, Illinois, USA

ABSTRACT *Staphylococcus epidermidis* is a native member of the human microbiota. Here, we report the draft genome sequence of *S. epidermidis* UMB8493, an isolate from a catheterized urine sample from a female with overactive bladder symptoms. The 2.54-Mbp draft genome encodes genes associated with beta-lactam and fosfomicin resistance.

Staphylococcus epidermidis is a Gram-positive, coagulase-negative staphylococcus (CoNS) traditionally considered to be a benign commensal microorganism found on the skin and mucous membranes of humans and other mammals (1). More recently, however, *S. epidermidis* has been identified as being the leading causative agent of and most prominently recovered *Staphylococcus* species from nosocomial infections associated with indwelling medical devices, including pacemakers and catheters (1, 2). *S. epidermidis* has routinely been isolated from urine samples (3). Prior research confirmed that it is present in the bladder microbiota and is not a contaminant from the skin microbiota during collection (4). Its role in the urinary microbiota remains unknown; it has not been conclusively linked with lower urinary tract symptoms in adults, although it has been reported to cause urinary tract infections in children (5, 6). Here, we present the draft genome sequence of *S. epidermidis* UMB8493, isolated from a urine specimen collected via a transurethral catheter from a woman with overactive bladder (OAB) symptoms.

S. epidermidis UMB8493 was isolated from a prior institutional review board (IRB)-approved study (Loyola University Chicago, approval number LU207102) using the expanded quantitative urinary culture (EQUC) method (3). This sample was collected from a patient at Loyola University Medical Center's Female Pelvic Medicine and Reconstructive Surgery Center (Maywood, IL, USA) in July 2018. The genus and species were determined using matrix-assisted laser desorption ionization–time of flight (MALDI-TOF) mass spectrometry following the previously described protocol (3) prior to storage at -80°C . From this freezer stock, the *S. epidermidis* isolate was streaked onto a Columbia nalidixic acid (CNA) agar plate and incubated at 35°C with 5% CO_2 for 24 h. A single colony was selected and inoculated in Trypticase soy broth and incubated at 37°C for 24 h. DNA was extracted using the DNeasy blood and tissue kit following the manufacturer's protocol for Gram-positive bacteria with the following exceptions: we used 230 μl of lysis buffer (180 μl of 20 mM Tris-Cl, 2 mM sodium EDTA, and 1.2% Triton X-100 and 50 μl of lysozyme) in step 2 and altered the incubation time in step 5 to 10 min. Upon purification, the genomic DNA was quantified using a Qubit fluorometer. DNA was then sent to the Microbial Genomic Sequencing Center (MiGS) at the University of Pittsburgh for sequencing. The libraries were prepared as follows: the DNA was first enzymatically fragmented using an Illumina tagmentation enzyme, and then

Citation Truckenbrod A, Miller-Ensminger T, Voukadinova A, Wolfe AJ, Putonti C. 2020. Draft genome sequence of *Staphylococcus epidermidis* UMB8493, isolated from the female urinary tract. *Microbiol Resour Announc* 9:e00419-20. <https://doi.org/10.1128/MRA.00419-20>.

Editor Julie C. Dunning Hotopp, University of Maryland School of Medicine

Copyright © 2020 Truckenbrod et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Catherine Putonti, cputonti@luc.edu.

Received 15 April 2020

Accepted 16 May 2020

Published 4 June 2020

indices were attached using PCR. The DNA library was sequenced using an Illumina NextSeq 550 platform, producing 2,584,013 pairs of 150-bp reads. Raw reads were trimmed using Sickle v1.33 (<https://github.com/najoshi/sickle>) and assembled using SPAdes v3.13.0 with the “only-assembler” option for K values of 55, 77, 99, and 127 (7). Genome coverage was calculated using BBDMap v38.47 (<https://sourceforge.net/projects/bbmap/>). The NCBI Prokaryotic Genome Annotation Pipeline (PGAP) v4.11 (8) and PATRIC (9) were used to annotate the genome sequence. Unless previously noted, default parameters were used for each software tool.

The *S. epidermidis* UMB8493 draft genome sequence is 2,535,022 bp long assembled into 61 contigs with a GC content of 32%, genome coverage of 267 \times , and N_{50} score of 88,403 bp. The PGAP annotation identified 2,316 protein-coding genes, 10 rRNAs (4 5S, 4 16S, and 2 23S), and 58 tRNAs. PATRIC detected several genes associated with antibiotic resistance, including *blaZ* (beta-lactam resistance) and *fosB* (fosfomycin resistance). Future analyses of *S. epidermidis* will further our understanding of the epidemiology and transmission of this important commensal microorganism in the female urinary microbiome, allowing us to develop more effective therapeutic interventions in the future.

Data availability. This whole-genome shotgun (WGS) project has been deposited in GenBank under the accession number [JAAUVT000000000](https://www.ncbi.nlm.nih.gov/GenBank/JAAUVT000000000). The version described in this paper is the first version, JAAUVT010000000. The raw sequencing reads have been deposited in the SRA under the accession number [SRR11441022](https://www.ncbi.nlm.nih.gov/SRA/SRR11441022).

ACKNOWLEDGMENTS

This work was conducted as part of Loyola University Chicago’s Department of Biology Bacterial Genomics course. For prior patient recruitment, we acknowledge the Loyola Urinary Education and Research Collaborative (LUEREC) and the patients who provided the samples for this study.

REFERENCES

- Otto M. 2009. Staphylococcus epidermidis: the “accidental” pathogen. *Nat Rev Microbiol* 7:555–567. <https://doi.org/10.1038/nrmicro2182>.
- Becker K, Heilmann C, Peters G. 2014. Coagulase-negative staphylococci. *Clin Microbiol Rev* 27:870–926. <https://doi.org/10.1128/CMR.00109-13>.
- Hilt EE, McKinley K, Pearce MM, Rosenfeld AB, Zilliox MJ, Mueller ER, Brubaker L, Gai X, Wolfe AJ, Schreckenberger PC. 2014. Urine is not sterile: use of enhanced urine culture techniques to detect resident bacterial flora in the adult female bladder. *J Clin Microbiol* 52:871–876. <https://doi.org/10.1128/JCM.02876-13>.
- Wolfe AJ, Toh E, Shibata N, Rong R, Kenton K, Fitzgerald M, Mueller ER, Schreckenberger P, Dong Q, Nelson DE, Brubaker L. 2012. Evidence of uncultivated bacteria in the adult female bladder. *J Clin Microbiol* 50:1376–1383. <https://doi.org/10.1128/JCM.05852-11>.
- Lozano V, Fernandez G, Spencer PL, Taylor SL, Hatch R. 2015. Staphylococcus epidermidis in urine is not always benign: a case report of pyelonephritis in a child. *J Am Board Fam Med* 28:151–153. <https://doi.org/10.3122/jabfm.2015.01.140118>.
- Upadhyayula S, Kambalapalli M, Asmar BI. 2012. Staphylococcus epidermidis urinary tract infection in an infant. *Case Rep Infect Dis* 2012:983153. <https://doi.org/10.1155/2012/983153>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
- Wattam AR, Davis JJ, Assaf R, Boisvert S, Brettin T, Bun C, Conrad N, Dietrich EM, Disz T, Gabbard JL, Gerdes S, Henry CS, Kenyon RW, Machi D, Mao C, Nordberg EK, Olsen GJ, Murphy-Olson DE, Olson R, Overbeek R, Parrello B, Pusch GD, Shukla M, Vonstein V, Warren A, Xia F, Yoo H, Stevens RL. 2017. Improvements to PATRIC, the all-bacterial Bioinformatics Database and Analysis Resource Center. *Nucleic Acids Res* 45:D535–D542. <https://doi.org/10.1093/nar/gkw1017>.