



5-11-2018

Caspase-3 suppresses diethylnitrosamine-induced hepatocyte death, compensatory proliferation and hepatocarcinogenesis through inhibiting p38 activation

Na Shang

Loyola University Chicago Stritch School of Medicine

Thomas Bank

Loyola University Chicago Stritch School of Medicine

Xianzhong Ding

Loyola University Chicago Stritch School of Medicine

Peter W. Breslin

Loyola University Chicago, pbreslin@luc.edu

Jun Li

University of Notre Dame
Follow this and additional works at: https://ecommons.luc.edu/biology_facpubs

 Part of the [Biology Commons](#)

See next page for additional authors

Recommended Citation

Shang, Na; Bank, Thomas; Ding, Xianzhong; Breslin, Peter W.; Li, Jun; Shi, Baomin; and Qiu, Wei. Caspase-3 suppresses diethylnitrosamine-induced hepatocyte death, compensatory proliferation and hepatocarcinogenesis through inhibiting p38 activation. *Cell Death & Disease*, 9, : , 2018. Retrieved from Loyola eCommons, Biology: Faculty Publications and Other Works, <http://dx.doi.org/10.1038/s41419-018-0617-7>

This Article is brought to you for free and open access by the Faculty Publications and Other Works by Department at Loyola eCommons. It has been accepted for inclusion in Biology: Faculty Publications and Other Works by an authorized administrator of Loyola eCommons. For more information, please contact ecommons@luc.edu.



This work is licensed under a [Creative Commons Attribution 4.0 International License](#).

© Shang et al., 2018.

Authors

Na Shang, Thomas Bank, Xianzhong Ding, Peter W. Breslin, Jun Li, Baomin Shi, and Wei Qiu

ARTICLE

Open Access

Caspase-3 suppresses diethylnitrosamine-induced hepatocyte death, compensatory proliferation and hepatocarcinogenesis through inhibiting p38 activation

Na Shang¹, Thomas Bank¹, Xianzhong Ding², Peter Breslin^{1,3}, Jun Li⁴, Baomin Shi⁵ and Wei Qiu^{1,5}

Abstract

It is critical to understand the molecular mechanisms of hepatocarcinogenesis in order to prevent or treat hepatocellular carcinoma (HCC). The development of HCC is commonly associated with hepatocyte death and compensatory proliferation. However, the role of Caspase-3, a key apoptotic executor, in hepatocarcinogenesis is unknown. In this study, we used *Caspase-3*-deficient mice to examine the role of Caspase-3 in hepatocarcinogenesis in a chemical (diethylnitrosamine, DEN)-induced HCC model. We found that *Caspase-3* deficiency significantly increased DEN-induced HCC. Unexpectedly, *Caspase-3* deficiency increased apoptosis induced by DEN and the subsequent compensatory proliferation. Intriguingly, we discovered that *Caspase-3* deficiency increased the activation of p38 with and without DEN treatment. Moreover, we demonstrated that TNF α and IL1 α stimulated increased activation of p38 in *Caspase-3* KO hepatocytes compared with wild-type hepatocytes. Finally, we found that inhibition of p38 by SB202190 abrogated enhanced hepatocyte death, compensatory proliferation and HCC induced by DEN in *Caspase-3*-deficient mice. Overall, our data suggest that Caspase-3 inhibits chemical-induced hepatocarcinogenesis by suppressing p38 activation and hepatocyte death.

Introduction

Hepatocellular carcinoma (HCC) is the second leading cause of cancer deaths worldwide¹. The overall survival of patients with HCC is <12%, and most patients with HCC have limited treatment options². Currently, the most effective targeted therapeutic agent for advanced HCC, sorafenib, only increases survival in patients with advanced HCC from 7.9 months to 10.7 months³. There is an urgent need to develop more effective therapeutic strategies and agents to treat HCC. To achieve this goal,

the molecular signaling pathways that drive or mediate the development of HCC must be better understood.

HCC is considered a chronic inflammation-related disorder, the development of which is commonly associated with hepatocyte death and compensatory proliferation^{1,4–10}. Several classes of chemicals promote HCC in rodents, including diethylnitrosamine (DEN), which has been extensively studied. In the DEN-induced HCC model, DEN stimulates DNA damage and causes reactive oxygen species to accumulate; it also induces hepatocyte death through the c-Jun N-terminal kinase (JNK) pathway^{11–14}. The dying cells release interleukin-1 alpha (IL1 α), which activates Kupffer cells^{6,15}. Activated Kupffer cells produce cytokines and growth factors such as interleukin-6 (IL-6) and tumor necrosis factor alpha (TNF α)^{5,16–18}, which promote expansion of hepatocytes with DNA mutations thus enhancing HCC

Correspondence: Baomin Shi (baomingsph@163.com) or Wei Qiu (wqiu@luc.edu)

¹Department of Surgery and Oncology Institute, Loyola University Chicago Stritch School of Medicine, 2160 South 1st Avenue, Maywood, IL 60153, USA

²Department of Pathology, Loyola University Chicago Stritch School of Medicine, 2160 South 1st Avenue, Maywood, IL 60153, USA

Full list of author information is available at the end of the article.

Edited by E. Baehrecke

© The Author(s) 2018



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.





Fig. 1 *Caspase-3* deficiency promoted DEN-induced liver cancer. **a** CASPASE-3, active CASPASE-3 and β -actin protein expression in the livers of WT mice 3 days following injection with either saline (Un) or 100 mg/kg of DEN was analyzed by western blotting. Values are means \pm SD, $n = 3$ mice in each group. **b** Active CASPASE-3 protein (red) in the livers of WT and *Caspase-3* KO mice following indicated treatment was detected by IF with nuclei counterstained with DAPI (magnification, $\times 400$). **c** Photographs of livers of WT and *Caspase-3* KO mice 9 months after DEN injection. **d** Quantification of liver tumor numbers ($n = 6$). **e** Quantification of liver tumor sizes ($n = 6$). **f** Histological analysis (H&E staining) and staining of AFP of the livers of WT and *Caspase-3* KO mice 9 months after DEN injection. Bars: 20 μ M. Values in **(d)** and **(e)** are means \pm SDs

development^{5,16,17}. We previously reported that deficiency of *PUMA* (p53 upregulated modulator of apoptosis) significantly decreased DEN-induced liver cancer by blocking acute apoptotic responses and the subsequent compensatory proliferation¹⁹. These data provide evidence for the correlation of specific mediators of hepatocyte apoptosis and the development of HCC. Caspase-3 is a member of the cysteine-aspartic acid protease family (caspases) and plays a central role in the execution-phase of cellular apoptosis²⁰. Given the strong association of apoptosis and compensatory proliferation in HCC development, and the specific apoptotic executive function of Caspase-3, we hypothesized that inhibition of Caspase-3

would suppress DNA damage-induced hepatocyte death, and thereby inhibit hepatic carcinogenesis.

Unexpectedly, we found that *Caspase-3* deficiency significantly promoted DEN-induced HCC development. Intriguingly, *Caspase-3* deficiency increased hepatocyte death and the subsequent compensatory proliferation induced by DEN. In addition, we discovered that *Caspase-3* deficiency increased the activation of p38 MAP kinase. Moreover, we demonstrated that deletion of *Caspase-3* increased p38 activation by IL1 α and TNF α in primary mouse hepatocytes. Furthermore, we found that inhibition of p38 abrogated enhanced hepatocyte death, the compensatory proliferation and HCCs induced by DEN in



Caspase-3-deficient mice. Overall, our data suggest that *Caspase-3* inhibits chemical-induced hepatocarcinogenesis by suppressing p38 activation and hepatocyte death.

Results

DEN induced *Caspase-3* cleavage in mouse livers

To determine the role of *Caspase-3* in hepatocarcinogenesis, we used the DEN-induced HCC model because *Caspase-3* is cleaved and activated in response to DNA

damage or non-genotoxic stimuli²¹ and DEN is known to induce DNA damage²². We analyzed the activation of *Caspase-3* in the livers of *WT* mice after DEN treatment. Cleaved *Caspase-3* but not total *Caspase-3* was significantly elevated by day 3 after treatment compared with untreated mice (Fig. 1a). Immunostaining indicated that cleaved *Caspase-3* protein was selectively induced in hepatocytes around the centrilobular regions 24 h after DEN treatment, whereas the basal level was undetectable (Fig. 1b).

Caspase-3 deficiency promoted DEN-induced liver cancer

A single injection of DEN to 15-day-old male mice results in efficient HCC induction¹⁹. To determine a potential role for Caspase-3 in hepatocarcinogenesis, we compared tumor incidence and size in *WT* and *Caspase-3* knockout (*Casp3* KO) littermates 9 months after DEN treatment. All of the mice developed tumors by 9 months of age (Fig. 1c). Interestingly, tumor incidence in *Casp3* KO mice increased by about twofold compared with *WT* mice (8.8 ± 2.2 vs. 3.5 ± 0.8) (Figs. 1c, d). Tumor size was also significantly increased in *Casp3* KO mice compared with *WT* mice (Figs. 1e, f). In addition, immunohistochemical (IHC) staining signals for alpha-fetoprotein (AFP), a common HCC marker, were significantly lower in the livers of *WT* mice compared with those of *Casp3* KO mice (Fig. 1f). These results indicate that *Casp3* deficiency promotes DEN-induced hepatocarcinogenesis.

Caspase-3 deficiency increased DEN-induced hepatocyte death

Caspase-3 is a critical apoptotic executor²¹. Promotion of HCC development by *Casp3* deficiency could be due to decreased apoptosis of tumor cells. To determine whether a deficiency of *Casp3* affects apoptosis in DEN-induced HCC, we performed Terminal deoxynucleotidyl transferase dUTP nick end labeling (TUNEL) staining in liver tumors from *WT* and *Casp3* KO mice treated with DEN. We did not find significant differences in apoptosis in tumors between these two groups of mice (Fig. S1A and S1B). We also analyzed proliferation in the DEN-treated livers from *WT* and *Casp3* KO mice by Ki67 staining. The number of Ki67⁺ cells was comparable in *Casp3*-deficient and *WT* liver tumors (Fig. S1A and S1B). These results show that *Casp3* deficiency affects neither apoptosis nor proliferation in DEN-induced HCC.

The development of DEN-induced HCC is associated with hepatocyte death and compensatory proliferation^{1,4–10}. DEN induces significant hepatocyte death in *WT* mice within 3 days (Figs. 2a, b), notably, in the centrilobular regions where Caspase-3 is activated (Figs. 1b, 2a). Interestingly, there was dramatically enhanced hepatocyte death in *Casp3* KO mice compared with *WT* mice after DEN treatment (Figs. 2a, b). Hepatocyte death decreased by 10 days after DEN treatment in both *WT* and *Casp3* KO mice, but was greater in *Casp3* KO mice than in *WT* mice (Figs. 2a, b). DEN treatment increased serum levels of the liver enzyme alanine aminotransferase (ALT) by day 3, such increase being similar in both *Casp3* KO and *WT* mice (Fig. 2c). As DEN can induce oxidative DNA damage in the liver⁵, we evaluated DNA double-strand breaks in livers by p-H2AX staining. No difference in DNA damage was found between *WT* and *Casp3* KO mice 3 or 10 days after DEN treatment (Figs. 2d, e).


Caspase-6 and Caspase-7 may play similar functions as Caspase-3 in the execution of apoptosis²³. To determine if deletion of *Caspase-3* increases the activation of Caspase-6 and Caspase-7 after DEN treatment in mouse livers, we examined the expression of total and cleaved Caspase-6 and Caspase-7 in DEN-treated mouse livers. We found a slight decrease in cleavage of Caspase-6 or Caspase-7 in *Casp3* KO mice compared with *WT* mice (Fig. S2), suggesting that increased DEN-induced hepatocyte apoptosis resulting from Caspase-3 deficiency is not the result of enhanced activation of Caspase-6 or Caspase-7 in DEN-treated mouse livers.

Caspase-3 deficiency enhanced DEN-induced compensatory proliferation

DEN-induced hepatocyte death is associated with compensatory proliferation and HCC development¹⁹. As expected, elevated proliferation was found in the livers of *WT* mice 3 and 10 days following DEN treatment as evidenced by Ki67 and PCNA staining (Figs. 3a–d). The proliferation centered on the centrilobular regions where apoptotic cells were detected (Figs. 2a, 3a). The degree of proliferation was significantly increased in *Casp3* KO mice compared with *WT* mice (Figs. 3a–d). IL-6 and its downstream target Stat3 have been shown to play important roles in DEN-induced compensatory proliferation^{5,16–18}. We therefore examined the expression of IL-6 mRNA and phosphorylation of Stat3 in *WT* and *Casp3* KO mouse livers. We found that IL-6 mRNA was increased in *Caspase-3* KO mouse livers compared with *WT* mouse livers after DEN treatment both on day 3 and day 10 (Fig. S3). Phosphorylation of Stat3 was also increased in *Caspase-3* KO mouse livers compared with *WT* mouse livers after DEN treatment on day 3 while the difference was diminished by day 10 (Fig. S3). These results showed that *Caspase3* deficiency promotes DEN-induced compensatory proliferation in hepatocytes.

Caspase-3 deficiency enhanced p38 activation in mouse livers

Activation of JNK, nuclear factor- κ B (NF- κ B) and p38 by cytokines, including TNF α and IL1 α , plays a critical role in hepatocyte death and compensatory proliferation after DEN treatment^{24,25}. To determine whether deletion of *Caspase-3* affects the activation of JNK, NF- κ B and p38 in the liver after DEN treatment, we compared the phosphorylation of these three proteins in the livers of *WT* and *Casp3* KO mice. There was no significant difference in the phosphorylation status of JNK in *Casp3* KO mouse livers compared with *WT* mouse livers prior to or after DEN treatment (Fig. 4a). NF- κ B was slightly increased by *Caspase-3* deficiency in response to DEN treatment by 3 days but not after 10 days (Fig. 4a). Accordingly, phosphorylation of I-kappa-B-alpha (I κ B α)




on Ser 32 and Ser 36, which results in release and nuclear translocation of active NF- κ B²⁶, was also slightly increased by *Caspase-3* deficiency in response to DEN treatment on day 3 after treatment (Fig. S4A). We also examined levels of I κ B α , Vcam1 and Cxcl10 mRNA, which have been reported to be direct target genes of NF- κ B^{27,28}. We found that the expression of Vcam1 and Cxcl10 but not I κ B α was increased by *Caspase-3* deficiency in response to DEN treatment in mouse livers (Fig. S4B). NF- κ B has been shown to protect against DEN-induced hepatocyte death and HCC^{5,24}. Therefore, it is unlikely that an increase in NF- κ B activation led to enhanced DEN-induced hepatocyte death, compensatory proliferation and HCC caused by *Caspase-3* deficiency. Interestingly, higher p-p38 was observed in *Casp3* KO mouse livers compared with *WT* mouse livers with or without DEN treatment (Fig. 4a), suggesting that deletion of *Caspase-3* increases the activation of p38 in mouse livers. Consistently, phosphorylation of MK2, a downstream direct target of p38, was also increased in *Casp3*

KO mouse livers (Fig. S5A). MMK6 and MMK3 have been shown to phosphorylate and activate p38. We also found that phosphorylation of MAP kinase kinases 3/6 (MKK3/6) was increased in *Casp3* KO mouse livers (Fig. S5A).

***Caspase-3* deficiency did not affect TNF α nor IL1 α expression in mouse livers after DEN treatment**

TNF α plays a critical role in hepatocyte compensatory proliferation and the activation of JNK, NF- κ B and p38 after DEN treatment^{24,25}. It is possible that *Caspase-3* deficiency enhances p38 activation in mouse liver by increasing the expression of TNF α after DEN treatment. We therefore compared mRNA and protein levels of TNF α in the livers of *WT* and *Casp3* KO mice. Expression of both TNF α mRNA and protein was increased when examined 3 days following DEN treatment in *WT* mice, whereas comparable expression of TNF α was observed in *Casp3* KO mice (Fig. 4b and Fig. S5A). IL1 α has been shown to phosphorylate and activate p38 in liver



cells²⁹. We therefore examined the expression of IL1 α and found no significant difference in *Il1 α* mRNA expression (Fig. 4b). These data indicate that *Caspase-3* deficiency increases the activation of p38 but does not do so through enhanced expression of TNF α or IL1 α in mouse livers.

Caspase-3 deficiency enhanced activation of p38 by in response to TNF α or IL1 α in hepatocytes

As the expression of neither TNF α nor IL1 α was affected in mouse livers by deletion of *Caspase-3*, we therefore examined whether *Caspase-3* deficiency in hepatocytes enhances the phosphorylation of p38 in

response to TNF α or IL1 α . We isolated primary hepatocytes from WT and *Casp3* KO mice and treated them with TNF α or IL1 α . There was a significant enhancement of p38 phosphorylation in *Casp3* KO hepatocytes compared with WT hepatocytes prior to or after treatment with TNF α (Fig. 4c). Similarly, phosphorylation of p38 was increased in *Casp3* KO hepatocytes compared with WT hepatocytes after 1 h of IL1 α treatment (Fig. 4d). Consistently, phosphorylation of MK2 and MKK3/6 was also increased in *Casp3* KO hepatocytes compared with WT hepatocytes after 1 h of treatment with IL1 α or TNF α (Fig. S5B and C). In general, these data demonstrate that *Caspase-3* deficiency enhances

the activation of p38 in hepatocytes in response to *TNF α* or *IL1 α* .

Inhibition of p38 abrogated enhanced DEN-induced hepatocyte death, compensatory proliferation and HCC development induced by deletion of Caspase-3

Emerging evidence indicates that activation of p38 promotes cell death. Activation of p38 by MKK6 has been shown to induce apoptosis in Huh7 and HepG2 cells³⁰. A p38 MAPK inhibitor markedly suppressed cell death induced by *TNF α* ³¹. Therefore, we reasoned that *Caspase-3* deficiency increased DEN-induced hepatocyte apoptosis through enhanced p38 activation. To test this hypothesis, we first examined whether the overexpression of p38 increases *TNF α* -induced cell death in AML12 cells, a mouse hepatocyte cell line. We found that overexpression of p38 did increase *TNF α* -induced cell death in these cells (Fig. S6). Further, we treated *WT* and *Casp3* KO mice with a specific inhibitor of p38, SB202190, prior to and after DEN treatment. We found that SB202190 decreased the phosphorylation of p38 in both *WT* and *Casp3* KO mice (Fig. 5a) and inhibited DEN-induced hepatocyte death (Figs. 5b, c). Consistently, SB202190 also decreased DEN-induced compensatory proliferation in *Casp3* KO mice (Fig. 5d). Importantly, SB202190 inhibited DEN-induced HCC development in *Casp3* KO mice (Figs. 5e, f). These data suggest that deletion of *Caspase-3* enhances DEN-induced hepatocyte death, compensatory proliferation and the development of HCC by increasing the activation of p38.

Discussion

Treatment options for HCC are limited². Currently, the most effective targeted therapeutic agent for advanced HCC, sorafenib, an inhibitor of several protein tyrosine kinases (including vascular endothelial growth factor receptor and platelet-derived growth factor receptor) and Raf kinases, increases survival in patients with advanced HCC by only 3 months³. More effective therapeutic strategies and agents to treat HCC are clearly needed. Therefore, it is necessary to elucidate the molecular signaling pathways that drive or mediate the development of HCC. Hepatocyte apoptosis has been shown to play an important role in HCC development. About 3.5% of HCC samples from The Cancer Genome Atlas (TCGA) liver HCC data sets show deletion of the *Caspase-3* gene, which is correlated with reduced *Caspase-3* mRNA expression (Fig. S7). A recent study indicated that low expression of *Caspase-3* is correlated with poor prognosis in HCC patients³², suggesting that *Caspase-3* might be involved in HCC pathogenesis. However, the role of *Caspase-3* in HCC development in vivo has not been reported. In this study, we found that deletion of *Caspase-3* promotes DEN-induced hepatocyte death,

compensatory proliferation and carcinogenesis, indicating that *Caspase-3* plays an inhibitory role in hepatocarcinogenesis. Therefore, it is likely that *Caspase-3* deletion or downregulation in human HCC samples promotes HCC development.

Caspase-3 is a central executor of apoptosis. Therefore, it was surprising to us that *Caspase-3* deletion increases apoptosis induced by DEN in mouse livers. *Caspase-6* and *Caspase-7*, the other two apoptotic executors, have been shown to compensate the functions of *Caspase-3* in some cell contexts³³. However, we observed no compensatory activation of *Caspase-6* or *Caspase-7* in DEN-treated *WT* or *Caspase-3* KO livers. Instead, we found that loss of *Caspase-3* increases the activation of p38 in response to DEN, *TNF α* or *IL1 α* in hepatocytes. It is notable that p38 hyperactivation in *Casp3* KO mice is largely constitutive (Fig. 4a), although treatment with *IL1 α* or *TNF α* promotes a further increase in phosphorylation (Fig. S5B). It is possible that *Caspase-3* deletion increases p38 activation in response to the basal level of *TNF α* or *IL1 α* in mouse liver. Activation of p38 by MKK6 has been shown to induce apoptosis in HCC cells³⁰. p38 also mediates cell death induced by *TNF α* in cancer cells³¹. We also found that overexpression of p38 enhanced *TNF α* -induced cell death in mouse liver cells (Fig. S6). In addition, we demonstrated that a p38 inhibitor abrogates the enhanced DEN-induced hepatocyte death induced by deletion of *Caspase-3*. Overall, our data suggest that *Casp3* KO hepatocytes are more sensitive to *TNF α* , which is likely due to p38 hyperactivation (Fig. S6). Therefore, deletion of *Caspase-3* enhances DEN-induced hepatocyte death, compensatory proliferation and hepatocarcinogenesis by increasing the activation of p38 (Fig. 6). To our knowledge, this is the first report of inhibition of p38 activation by *Casp-3*. It is unknown whether this mechanism is a general one that will be found all in cell types or whether it is in cell context specific. It will be interesting for us to test this question in all our future studies.

Increased activation of p38 caused by the deletion of *Caspase-3* may be due to the result of multiple actions. MKKs 3 and 6 and have been identified as common activators of p38 in the liver³⁴. Indeed we found that phosphorylation of MKK6/3 is increased in *Caspase-3*-deficient hepatocytes (Fig. S4), suggesting that *Caspase-3* deletion increases the activation of p38 through enhancing MKK6/3 activation in hepatocytes. Several MAP kinase kinase kinases have been shown to trigger MKK6/3 activation; they include apoptosis signal-regulating kinase 1 (ASK1), dual-leucine zipper-bearing kinase 1 (DLK1), thousand-and-one amino acid (TAO) 1 and 2, tumor progression loci 2 (TPL2), mixed lineage kinase 3 (MLK3), MEK kinase 3 (MEKK3), MEKK4, and leucine zipper and sterile- α motif kinase 1 (ZAK1)²⁹. Over three hundred targets of *Caspase-3* have been identified³⁵. It is





Fig. 5 Inhibition of p38 abrogated enhanced DEN-induced hepatocyte death, compensatory proliferation and HCC development induced by deletion of *Caspase-3*. **a** Expression of p-p38 and GAPDH proteins in the livers of *Caspase-3* KO mice 3 days following injection with either saline (Un) or 100 mg/kg DEN plus 20 mg/kg SB202190 was analyzed by western blotting. **b** Apoptosis in the livers of *Caspase-3* KO mice 3 days following injection with either saline (Un) or 100 mg/kg DEN plus 20 mg/kg SB202190 was analyzed by TUNEL staining. Bars: 20 μ M. **c** Quantification of TUNEL staining for **(b)** ($n = 3$). **d** Quantification of Ki67 staining in the livers of *Caspase-3* KO mice 3 days after injection with either saline (Un) or 100 mg/kg DEN plus 20 mg/kg SB202190. **e** Photographs of livers of *WT* and *Caspase-3* KO mice 9 months after DEN injection with or without SB202190. **f** Quantification of liver tumor numbers ($n = 5$) and liver tumor sizes ($n = 5$) for **(e)**. Values in **(c)**, **(d)** and **(f)** are means \pm SDs

possible that the loss of *Caspase-3* increases the activation of MKK6/3 and p38 through regulating its downstream targets. Given the cell context dependence and

multiple known targets of *Caspase-3*, it is not our contention that the increased activation of MKK6/p38 is the only mechanism to explain *Caspase-3*'s effects. Rather,



we posit that this mechanism is a contributor to increased hepatocyte death, compensatory proliferation and hepatic carcinogenesis caused by the loss of *Caspase-3*, and can account for some, if not all, of the functional changes observed.

In conclusion, our study shows that deficiency of *Caspase-3* promotes chemical-induced hepatocyte compensatory proliferation and hepatocarcinogenesis. Recent studies suggest that specific inhibition of Caspase-3 in concert with chemotherapy may be a novel approach for the treatment of cancers because of the role of cell death in promoting tissue regeneration in mammalian cells^{36,37}. Our data suggest that the role of Caspase-3 in cancer development is complex and the inhibition of Caspase-3 might not be suitable for the treatment of HCC patients.

Materials and methods

Mice and treatments

All animals received humane care according to the “Guide for the Care and Use of Laboratory Animals” (http://oacu.od.nih.gov/ac_cbt/guide3.htm). The procedures for all animal experiments were approved by the Institutional Animal Care and Use Committee of Loyola University Chicago. *Casp3*^{+/+} and *Casp3*^{-/-} littermate mice in a C57BL/6 background were generated from heterozygote intercrosses. Genotyping was performed as described in the JAX Lab

website (<https://www.jax.org/strain/006233>). The mice were housed in micro-isolator cages in a room illuminated from 07:00 a.m. to 07:00 p.m. (12:12-h light–dark cycle), and allowed access to water and chow ad libitum.

For the DEN-induced HCC model, DEN (15 mg/kg) was injected intraperitoneally (i.p.) into 15-day-old mice. Mice were sacrificed after 9 months on the standard diet. Surface tumor nodules in each liver lobe were counted and measured with a caliper. For short-term studies of DEN-induced liver injury, 8- to 12-week-old mice were injected with DEN (100 mg/kg body weight) i.p. and sacrificed after 3 or 10 days.

For p38 inhibitor treatments, SB202190 (LC Laboratories, Cat # S-1700) was diluted in 6% captisol (CyDex, Inc., Lenexa, KS). One day prior to DEN treatment, six 2-week-old or 8- to 12-week-old WT and six 2-week-old or 8- to 12-week-old *Casp3*^{-/-} mice were administered vehicle solution (6% captisol) or 20 mg/kg SB202190 by i. p. daily for 3 days and livers were collected 9 months, 3 or 10 days after DEN treatment.

Isolation and culture of primary mouse hepatocytes

Hepatocytes from *Casp3*^{+/+} and *Casp3*^{-/-} mice were isolated by the non-recirculating two-step perfusion method as previously described³⁸. The hepatocytes were then cultured in Williams’ medium E supplemented with Hepatocyte Maintenance Supplement Pack (Life Technologies, Grand Island, NY) overnight. Cells were treated with 5 ng/ml and 20 ng/ml IL1 α or 5 ng/ml and 15 ng/ml TNF α for 1 h; proteins were subsequently collected for western blot analysis.

Culture and treatment of AML12 cells

AML12 cells were purchased from ATCC and cultured as per the manufacturer’s recommendations. Cells were transfected with pCNA3 or pCNA3-p38 α (Addgene, #20352) plasmids followed by treatment with either saline (vehicle) or 20 ng/ml TNF α for 48 h. Viability of the cells was analyzed by Alamar Blue Assay.

ALT measurement

Venous blood of mice was collected from the tail vein 0, 24, and 72 h after DEN treatment. Blood was kept at 4 °C for overnight and centrifuged at 200 \times g for 20 min to isolate serum. ALT was measured using the Infinity™ ALT kit (Thermo Scientific, Middletown, VA) and is reported as mean \pm SD. Briefly, 10 μ l of serum was added to 100 μ l ALT reagent and measured at 37 °C at 340 nm. Each sample was measured in triplicate and three mice were used in each group.

Western blotting

Western blotting was performed as previously described³⁸. Primary antibodies, against Caspase-3 (9662), active

Caspase-3 (9661), Caspase-6 (9762), active Caspase-6 (9761), Caspase-7 (9492), active Caspase-7 (8438), JNK (9252), p-JNK (9255), NF- κ B (8242), p-NF- κ B (3033), p38 (8690), p-p38 (4631), p-MK2 (3007), MKK3 (8535), MKK6 (8550), p-MKK3/6 (12280), were purchased from Cell Signaling Technologies (Danvers, MA).

TUNEL staining

TUNEL staining was performed as previously described^{38,39}. The apoptotic index was scored in at least five fields at $\times 400$ magnification/mouse and reported as mean \pm SD. Three mice were used for each group.

IHC staining

IHC was performed as previously described^{38,40}. Cells with positive staining were scored in at least five fields at $\times 400$ or $\times 200$ magnification and reported as mean \pm SD. Three mice were used per group.

Total RNA extraction and real-time reverse transcriptase polymerase chain reaction

Approximately 100 mg of fresh tissue was minced and put into 600 μ l lysis buffer (Promega). Total RNA was isolated, and complementary DNA was then generated for real-time PCR analysis as described⁴⁰. Real-time PCR was performed on a Mini Opticon Real-time PCR system (Bio-Rad) with SYBR Green (Invitrogen) and specific primers¹⁹. Melting curves and agarose gel electrophoresis of the PCR products were used to verify the specificity of PCR amplification.

Statistical analysis

Statistical analysis was performed using GraphPad Prism V software. Data are presented as means \pm standard deviations (SD). Statistical significance was calculated using the Student's *t*-test. $P < 0.05$ was considered to be significant. The means \pm SDs are shown in figures where applicable.

Acknowledgements

We thank Drs. Mitchell F. Denning, Nancy Zeleznik-Le, Jiwang Zhang and Manuel Diaz (all of Loyola University Chicago) for their helpful discussions and advice. This work is supported in part by AASLD Liver Scholar Award (W.Q.), NIH R03CA195183 (W.Q.), R03CA184652 (W.Q.) and R01CA197128 (W.Q.).

Author details

¹Department of Surgery and Oncology Institute, Loyola University Chicago Stritch School of Medicine, 2160 South 1st Avenue, Maywood, IL 60153, USA. ²Department of Pathology, Loyola University Chicago Stritch School of Medicine, 2160 South 1st Avenue, Maywood, IL 60153, USA. ³Department of Molecular/Cellular Physiology, Loyola University Chicago Stritch School of Medicine, 2160 South 1st Avenue, Maywood, IL 60153, USA. ⁴Department of Applied and Computational Mathematics and Statistics, University of Notre Dame, Notre Dame, IN, USA. ⁵Department of General Surgery Tongji Hospital, Tongji University Medical School, Shanghai, PR 200065, China

Authors' contributions

N.S. performed experiments, analyzed data and wrote the paper. T.B. bred mice and performed experiments. P.B. wrote the paper. X.Z.D. analyzed histology. J.L. performed statistical analysis. B.M.S. designed experiments, analyzed data and wrote the paper. W.Q. designed and performed experiments, analyzed data and wrote the paper.

Conflict of interest

The authors declare that they have no conflict of interest.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Supplementary Information accompanies this paper at (<https://doi.org/10.1038/s41419-018-0617-7>).

Received: 12 April 2018 Accepted: 19 April 2018

Published online: 11 May 2018

References

- Bruix, J. & Sherman, M., Practice Guidelines Committee, AA.f.t.S.o.L.D. Management of hepatocellular carcinoma. *Hepatology* **42**, 1208–1236 (2005).
- Johnson, P. J. Non-surgical treatment of hepatocellular carcinoma. *HPB* **7**, 50–55 (2005).
- Llovet, J. M. et al. Sorafenib in advanced hepatocellular carcinoma. *N. Engl. J. Med.* **359**, 378–390 (2008).
- Bruix, J., Boix, L., Sala, M. & Llovet, J. M. Focus on hepatocellular carcinoma. *Cancer Cell.* **5**, 215–219 (2004).
- Maeda, S., Kamata, H., Luo, J. L., Leffert, H. & Karin, M. IKKbeta couples hepatocyte death to cytokine-driven compensatory proliferation that promotes chemical hepatocarcinogenesis. *Cell* **121**, 977–990 (2005).
- He, G. & Karin, M. NF-kappaB and STAT3 - key players in liver inflammation and cancer. *Cell Res.* **21**, 159–168 (2011).
- Schulte-Hermann, R., Bursch, W., Low-Baselli, A., Wagner, A. & Grasl-Kraupp, B. Apoptosis in the liver and its role in hepatocarcinogenesis. *Cell. Biol. Toxicol.* **13**, 339–348 (1997).
- Farber, E. Hepatocyte proliferation in stepwise development of experimental liver cell cancer. *Dig. Dis. Sci.* **36**, 973–978 (1991).
- Aravalli, R. N., Cressman, E. N. & Steer, C. J. Cellular and molecular mechanisms of hepatocellular carcinoma: an update. *Arch. Toxicol.* **87**, 227–247 (2012).
- Weber, A., Boege, Y., Reisinger, F. & Heikenwalder, M. Chronic liver inflammation and hepatocellular carcinoma: persistence matters. *Swiss Med. Wkly.* **141**, w13197 (2011).
- Sakurai, T., Maeda, S., Chang, L. & Karin, M. Loss of hepatic NF-kappa B activity enhances chemical hepatocarcinogenesis through sustained c-Jun N-terminal kinase 1 activation. *Proc. Natl. Acad. Sci. USA* **103**, 10544–10551 (2006).
- Schwabe, R. F. et al. c-Jun-N-terminal kinase drives cyclin D1 expression and proliferation during liver regeneration. *Hepatology* **37**, 824–832 (2003).
- Hui, L., Zatloukal, K., Scheuch, H., Stepniak, E. & Wagner, E. F. Proliferation of human HCC cells and chemically induced mouse liver cancers requires JNK1-dependent p21 downregulation. *J. Clin. Invest.* **118**, 3943–3953 (2008).
- Seki, E., Brenner, D. A. & Karin, M. A liver full of JNK: signaling in regulation of cell function and disease pathogenesis, and clinical approaches. *Gastroenterology* **143**, 307–320 (2012).
- Sakurai, T. et al. Hepatocyte necrosis induced by oxidative stress and IL-1 alpha release mediate carcinogen-induced compensatory proliferation and liver tumorigenesis. *Cancer Cell.* **14**, 156–165 (2008).
- Naugler, W. E. et al. Gender disparity in liver cancer due to sex differences in MyD88-dependent IL-6 production. *Science* **317**, 121–124 (2007).
- Schwabe, R. F. & Brenner, D. A. Mechanisms of Liver Injury. I. TNF-alpha-induced liver injury: role of IKK, JNK, and ROS pathways. *Am. J. Physiol. Gastrointest. Liver Physiol.* **290**, G583–G589 (2006).
- Johnson, C. et al. Interleukin-6 and its receptor, key players in hepatobiliary inflammation and cancer. *Transl. Gastrointest. Cancer* **1**, 58–70 (2012).
- Qiu, W. et al. PUMA-mediated apoptosis drives chemical hepatocarcinogenesis in mice. *Hepatology* **54**, 1249–1258 (2011).

20. Crawford, E. D. & Wells, J. A. Caspase substrates and cellular remodeling. *Annu. Rev. Biochem.* **80**, 1055–1087 (2011).
21. Parrish, A. B., Freel, C. D. & Kornbluth, S. Cellular mechanisms controlling caspase activation and function. *Cold Spring Harb. Perspect. Biol.* **5**, 1–25 (2013).
22. Feinberg, A. & Zedeck, M. S. Production of a highly reactive alkylating agent from the organospecific carcinogen methylazoxymethanol by alcohol dehydrogenase. *Cancer Res.* **40**, 4446–4450 (1980).
23. Shi, Y. Caspase activation, inhibition, and reactivation: a mechanistic view. *Protein Sci.* **13**, 1979–1987 (2004).
24. Karin, M. NF-kappaB as a critical link between inflammation and cancer. *Cold Spring Harb. Perspect. Biol.* **1**, a000141 (2009).
25. Breitwieser, W. et al. Feedback regulation of p38 activity via ATF2 is essential for survival of embryonic liver cells. *Genes Dev.* **21**, 2069–2082 (2007).
26. Chen, Z. J., Parent, L. & Maniatis, T. Site-specific phosphorylation of Ikkappa-Balpha by a novel ubiquitination-dependent protein kinase activity. *Cell* **84**, 853–862 (1996).
27. Hein, H. et al. Genomic organization, sequence, and transcriptional regulation of the human eotaxin gene. *Biochem. Biophys. Res. Commun.* **237**, 537–542 (1997).
28. Pahl, H. L. Activators and target genes of Rel/NF-kappaB transcription factors. *Oncogene* **18**, 6853–6866 (1999).
29. Koul, H. K., Pal, M. & Koul, S. Role of p38 MAP kinase signal transduction in solid tumors. *Genes Cancer* **4**, 342–359 (2013).
30. Iyoda, K. et al. Involvement of the p38 mitogen-activated protein kinase cascade in hepatocellular carcinoma. *Cancer* **97**, 3017–3026 (2003).
31. Kang, J. S. et al. DBM1285 suppresses tumor necrosis factor alpha production by blocking p38 mitogen-activated protein kinase/mitogen-activated protein kinase-activated protein kinase 2 signaling pathway. *J. Pharmacol. Exp. Ther.* **334**, 657–664 (2010).
32. Huang, H. et al. Expression and prognostic significance of osteopontin and caspase-3 in hepatocellular carcinoma patients after curative resection. *Cancer Sci.* **101**, 1314–1319 (2010).
33. McIlwain, D. R., Berger, T. & Mak, T. W. Caspase functions in cell death and disease. *Cold Spring Harb. Perspect. Biol.* **5**, a008656 (2013).
34. Enslen, H., Raingeaud, J. & Davis, R. J. Selective activation of p38 mitogen-activated protein (MAP) kinase isoforms by the MAP kinase kinases MKK3 and MKK6. *J. Biol. Chem.* **273**, 1741–1748 (1998).
35. Fischer, U., Janicke, R. U. & Schulze-Osthoff, K. Many cuts to ruin: a comprehensive update of caspase substrates. *Cell Death Differ.* **10**, 76–100 (2003).
36. Li, F. et al. Apoptotic cells activate the “phoenix rising” pathway to promote wound healing and tissue regeneration. *Sci. Signal.* **3**, ra13 (2010).
37. Boland, K., Flanagan, L. & Prehn, J. H. Paracrine control of tissue regeneration and cell proliferation by Caspase-3. *Cell Death Dis.* **4**, e275 (2013).
38. Shang, N. et al. FAK is required for c-Met/beta-catenin-driven hepatocarcinogenesis. *Hepatology* **61**, 214–226 (2015).
39. Shang, N. et al. FAK kinase activity is required for the progression of c-Met/beta-catenin-driven HCC. *Gene Exp.* **17**, 79–88 (2016).
40. Arteaga, M. et al. Inhibition of SIRT2 suppresses hepatic fibrosis. *Am. J. Physiol. Gastrointest. Liver. Physiol.* **310**, G1155–G1168 (2016).