

Expression of mRNA of the *col2a1b* gene during Zebrafish Embryogenesis

Mansi V. Patel, Mia S. Gabelev, Alex Tollefson, Arturo G. Gutiu, Vanessa L. Pahlow, Timoteea A. Saitis, Austin P. Runde and Rodney M. Dale

Loyola University Chicago, Department of Biology, Chicago, IL 60660



Abstract

Cartilage is an important tissue for all vertebrate organisms; it allows for flexibility yet structure in the extracellular matrix, which translates to support across nearly all parts of the body. Most cartilage elements are composed of two major cell types: chondrocytes and perichondrial cells. The perichondrium plays an important role in signaling to chondrocytes during the conversion of cartilage to bone (ossification). Mutations in genes/proteins expressed in these cell types can lead to morbidities such as osteoarthritis, Czech dysplasia, & Langer-Saldino achondroplasia, among others.

Our lab is interested in characterizing the genetic regulation of the critical structural genes that allow for the proper formation of cartilage. We set out to characterize the expression of *col2a1b*, the zebrafish ortholog of the human *COL2A1*. In zebrafish, the *col2a1a* gene (homolog of *col2a1b*) has been shown to be expressed in both chondrocytes and perichondrial cells, our preliminary work suggests *col2a1b* is only expressed in the perichondrium of cartilage elements; thus, the protein product of *col2a1b* is a type II alpha 1 collagen chain, which promotes cartilage development and ossification of chondrocytes. Our lab is interested in understanding the differential expression of the two homologs. In this report, we document the complete spatiotemporal expression of the zebrafish *col2a1b* mRNA during the first 5 days of development via *in situ* hybridization.

Danio rerio




Figure 1: Representative example of our model organism, the zebrafish – *Danio rerio*.

In Situ Hybridization (ISH) Mechanism




Figure 2: Schematic of the *In Situ* Hybridization technique. *In situ* hybridization (ISH) is a technique which allows for the sensitive and specific localization of mRNA fragments within a model organism, ours being the zebrafish. ISH relies on the principles of Watson-Crick nucleic acid complementarity, with hybridization ideally occurring between the endogenous mRNA strand and our lab-engineered, enzyme-conjugated, complementary ribonucleotide probe. *In situ* hybridization is an excellent method to use when attempting to characterize the spatiotemporal expression of genes critical to an organism's successful embryogenesis.

In Situ Hybridization of mRNA of *col2a1b*




Figure 4: Expression of *col2a1b* during days 1, 2, 3, 4, 5 post-fertilization.

Post-ISH, Light & Fluorescent Microscopy




Figure 5: Ventral view of expression during day 2 & 3 post fertilization. A. Two-day B. Three-day old zebrafish embryo showing craniofacial cartilage. C. Location of cartilage elements at 3 dpf fish. Modified from Brunt, *et al.* 2016.

Gel Electrophoresis of RT-PCR Product




Figure 3: RT-PCR of *col2a1b* mRNA extracted during development.

Results

In situ hybridization provided a baseline to compare endogenous gene expression to regulation elements reporter plasmids. From this we can determine the region in the genome responsible for *col2a1b* perichondrial expression. We have seen variable results after performing multiple *in situ* hybridizations, so we have made adjustments to ensure more consistent results in the future.

Future Directions

1. Cryosections of whole-mount zebrafish will be performed to determine more specific tissues of staining.
2. Reverse Transcription Polymerase Chain Reaction (RT-PCR) will be performed on all stages from 4-cell to 5 dpf.
3. Establish a system to allow for the repopulation of genetic lines.

Acknowledgments

Special thanks to Dr. Rodney M. Dale for being our mentor throughout our research at Loyola University Chicago; as well, thanks to the other members of the Dale Lab for collaborating efficiently and constructively. This research project was funded by the LUC *Mulcahy Scholars Program*.

References

1. Brunt, Lucy H., Norton, Joanna L., Bright, Jen A., Rayfield, Emily J., Hammond, Chrissy L. "Finite element modelling predicts changes in joint shape and cell behaviour due to loss of muscle strain in jaw development" *Journal of Biomechanics* 48 3112-3122. (2015)
2. Dale, R.M., and Topczewski, J. "Identification of an Evolutionarily Conserved Regulatory Element of the Zebrafish Collagen 2 Alpha 1a Gene." *Developmental Biology*. 357:518-531 (2011);
3. Haffner, P., Granato, M., Brand, M., Mullins, M.C., Hammerschmidt, Kane, D.A., Odenthal, J., van Eeden, F.J.M., Jiang, Y., Heisenberg, C., Kelsh, R.N., Furutani-Seiki, M., Vogelsang, E., Beuchle, D., Schach, U., Fabian, C., and Nüsslein-Volhard, C. "The identification of genes with unique and essential functions in the development of the zebrafish, *Danio rerio*." *Development*. 123:1-6. (1996)
4. Rodrigues, M., Lengerer, B., Ostermann, T., and Ladurner, P. "Molecular Biology Approaches in Bioadhesion Research." *Beilstein Journal of Nanotechnology*. 5:983-993. (2014)
5. Thisse, B., and Thisse C. "*In Situ* Hybridization on Whole-Mount Zebrafish Embryos and Young Larvae." *In situ hybridization protocols*. 1211:53-67. (2014)