



7-2017

Draft Genome Sequence of *Micrococcus luteus* (Schroeter) Cohn (ATCC 12698)

Catherine Putonti

Loyola University Chicago, cputonti@luc.edu

Evan Cudone

Loyola University Chicago

Laurynas Kalesinskas

Loyola University Chicago

Kathleen C. Engelbrecht

David W. Koenig

See next page for additional authors

Recommended Citation

Putonti, Catherine; Cudone, Evan; Kalesinskas, Laurynas; Engelbrecht, Kathleen C.; Koenig, David W.; and Wolfe, Alan J.. Draft Genome Sequence of *Micrococcus luteus* (Schroeter) Cohn (ATCC 12698). *Genome Announcements*, 5, 27: , 2017. Retrieved from Loyola eCommons, Biology: Faculty Publications and Other Works, <http://dx.doi.org/10.1128/genomeA.00576-17>

This Article is brought to you for free and open access by the Faculty Publications at Loyola eCommons. It has been accepted for inclusion in Biology: Faculty Publications and Other Works by an authorized administrator of Loyola eCommons. For more information, please contact ecommons@luc.edu.



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

© 2017 Putonti et al.

Authors

Catherine Putonti, Evan Cudone, Laurynas Kalesinskas, Kathleen C. Engelbrecht, David W. Koenig, and Alan J. Wolfe



Draft Genome Sequence of *Micrococcus luteus* (Schroeter) Cohn (ATCC 12698)

 Catherine Putonti,^{a,b,c,d} Evan Cudone,^a Laurynas Kalesinskas,^{a,b} Kathleen C. Engelbrecht,^e David W. Koenig,^e Alan J. Wolfe^d

Bioinformatics Program, Loyola University Chicago, Chicago, Illinois, USA^a; Department of Biology, Loyola University Chicago, Chicago, Illinois, USA^b; Department of Computer Science, Loyola University Chicago, Chicago, Illinois, USA^c; Department of Microbiology and Immunology, Stritch School of Medicine, Health Sciences Division, Loyola University Chicago, Maywood, Illinois, USA^d; Corporate Research and Engineering, Kimberly-Clark Corporation, Neenah, Wisconsin, USA^e

ABSTRACT The actinobacterium *Micrococcus luteus* can be found in a wide variety of habitats. Here, we report the 2,411,958-bp draft genome sequence of the type strain *M. luteus* (Schroeter) Cohn (ATCC 12698). Characteristic of this taxa, the genome sequence has a high G+C content, 73.14%.

Strains of the actinobacterium *Micrococcus luteus* have been isolated from a wide variety of habitats, including soil (1) and the microbiota of mammalian skin (2). Genomic sequences derived from these different sources provide insight into the diversity of this species (2–4). Here, we report the genome sequence and annotation of the type strain *Micrococcus luteus* (Schroeter) Cohn (ATCC 12698).

The culture isolate was purchased from ATCC and grown on 5% sheep blood agar (BD BBL prepared plated media) under 5% CO₂ for 48 h at 35°C. To extract genomic DNA, cells were resuspended in 0.5 mL of DNA extraction buffer (20 mM Tris-Cl, 2 mM EDTA, 1.2% Triton X-100, pH 8), followed by the addition of 50 μL of lysozyme (20 mg/mL), 30 μL of mutanolysin, and 5 μL of RNase (10 mg/mL). After a 1-h incubation at 37°C, 80 μL of 10% SDS and 20 μL of proteinase K were added. This mixture was then incubated for 2 h at 55°C. Then, 210 μL of 6M NaCl and 700 μL of phenol-chloroform were added. After a 30-min incubation with rotation, the solutions were centrifuged at 13,500 rpm for 10 min, and the aqueous phase was extracted. An equivalent volume of isopropanol was added and incubated for 10 min prior to centrifugation at 13,500 rpm for 10 min. The supernatant was decanted, and the DNA pellet was precipitated using 600 μL of 70% ethanol. Following ethanol evaporation, the DNA pellet was resuspended in Tris-EDTA (TE) and stored at –20°C.

Genomic DNA was diluted in water to a concentration of 0.2 ng/μL, as measured by a fluorometric-based method (Life Technologies, Inc.). Library preparation of 1 ng (5 μL) of isolated DNA was performed using the Nextera XT DNA library preparation kit. The library was sequenced on the MiSeq sequencer (Illumina) using the MiSeq version 2 reagent kit (500 cycles), and 1,645,841 paired-end reads were produced. Reads were first processed, removing adapter sequences and phiX contaminants, using BBDuk from the BBMap package (<http://sourceforge.net/projects/bbmap>). Trimmed reads were assembled using SPAdes version 3.5 (5) and scaffolded using SSPACE (6). This resulted in 126 contigs at an average coverage of 268.7×. Contigs varied in length from 1,181 bp to 149,560 bp (N_{50} = 31,560 bp). The total genome size was 2,411,958 bp with an observed G+C content of 73.14%. Annotations were produced using the software tool Peasant (7). Five rRNAs, 48 tRNAs, and 2,317 protein-coding sequences were detected. No regularly interspaced short palindromic repeat sequences (CRISPRs) were found within the *M. luteus* strain Cohn (ATCC 12698) genomic sequence (8); to date, the CRISPR/Cas system has not been identified in this taxon (9). The assembly for the ATCC

Received 14 May 2017 Accepted 16 May 2017 Published 6 July 2017

Citation Putonti C, Cudone E, Kalesinskas L, Engelbrecht KC, Koenig DW, Wolfe AJ. 2017. Draft genome sequence of *Micrococcus luteus* (Schroeter) Cohn (ATCC 12698). Genome Announc 5:e00576-17. <https://doi.org/10.1128/genomeA.00576-17>.

Copyright © 2017 Putonti et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Catherine Putonti, cputonti@luc.edu.

12698 strain and the representative genome for the strain *M. luteus* NCTC 2665 (GenBank no. NC_012803.1) were compared using Mauve (10), revealing polymorphisms as well as regions unique to each strain.

Accession number(s). The draft whole-genome project for *M. luteus* strain Cohn (ATCC 12698) has been deposited at DDBJ/EMBL/GenBank under the accession number [NAQZ00000000](https://doi.org/10.1093/bioinformatics/btq683). Raw sequence reads were deposited at DDBJ/EMBL/GenBank under the accession number [SRR5364946](https://doi.org/10.1093/bioinformatics/btq683).

ACKNOWLEDGMENTS

We acknowledge Gina Kuffel and Michael Zilliox for sequencing this genome and David Paulucki's assistance in culturing the strain. This work was supported by Kimberly-Clark Worldwide, Inc.

REFERENCES

1. Sims GK, Sommers LE, Konopka A. 1986. Degradation of pyridine by *Micrococcus luteus* isolated from soil. *Appl Environ Microbiol* 51: 963–968.
2. Young M, Artsatbanov V, Beller HR, Chandra G, Chater KF, Dover LG, Goh EB, Kahan T, Kaprelyants AS, Kyrpides N, Lapidus A, Lowry SR, Lykidis A, Mahillon J, Markowitz V, Mavromatis K, Mukamolova GV, Oren A, Rokem JS, Smith MCM, Young DI, Greenblatt CL. 2010. Genome sequence of the Fleming strain of *Micrococcus luteus*, a simple free-living actinobacterium. *J Bacteriol* 192:841–860. <https://doi.org/10.1128/JB.01254-09>.
3. Rakhshiya PM, Patel PP, Thaker VS. 2015. Whole genome sequences and annotation of *Micrococcus luteus* SUBG006, a novel phytopathogen of mango. *Genom Data* 6:10–11. <https://doi.org/10.1016/j.gdata.2015.07.022>.
4. Hanafy RA, Couger MB, Baker K, Murphy C, O'Kane SD, Budd C, French DP, Hoff WD, Youssef N. 2016. Draft genome sequence of *Micrococcus luteus* strain O'Kane implicates metabolic versatility and the potential to degrade polyhydroxybutyrates. *Genom Data* 9:148–153. <https://doi.org/10.1016/j.gdata.2016.08.006>.
5. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
6. Boetzer M, Henkel CV, Jansen HJ, Butler D, Pirovano W. 2011. Scaffolding pre-assembled contigs using SSPACE. *Bioinformatics* 27:578–579. <https://doi.org/10.1093/bioinformatics/btq683>.
7. Brenner J, Kalesinskas L, Putonti C. 2017. Exploring the diversity of *Bacillus* whole genome sequencing projects using Peasant, the prokaryotic assembly and annotation tool. *bioRxiv* 132084. <https://doi.org/10.1101/132084>.
8. Grissa I, Vergnaud G, Pourcel C. 2007. CRISPRFinder: a web tool to identify clustered regularly interspaced short palindromic repeats. *Nucleic Acids Res* 35:W52–W57. <https://doi.org/10.1093/nar/gkm360>.
9. Grissa I, Vergnaud G, Pourcel C. 2007. The CRISPRdb database and tools to display CRISPRs and to generate dictionaries of spacers and repeats. *BMC Bioinformatics* 8:172. <https://doi.org/10.1186/1471-2105-8-172>.
10. Darling ACE, Mau B, Blattner FR, Perna NT. 2004. Mauve: multiple alignment of conserved genomic sequence with rearrangements. *Genome Res* 14:1394–1403. <https://doi.org/10.1101/gr.2289704>.